
Status: Created

Doc ID: GEN02-0223.1

Effective Date: 02/12/2023

Last Review Date: 09/21/2022

Approval and implementation dates for specific health plans may vary. Please consult the applicable health plan for more details.

Clinical Appropriateness Guidelines

Genetic Testing

Appropriate Use Criteria: Somatic Tumor Testing

Proprietary

© 2023 Carelon Medical Benefits Management, Inc. All rights reserved.

Table of Contents

Description and Application of the Guidelines	3
General Clinical Guideline	4
Somatic Testing of Solid Tumors.....	6
Clinical Indications	6
General Requirements	6
Umbrella Criteria	6
Somatic Genomic Testing (Tumor Biomarker Testing).....	6
Metastatic or Advanced Cancer (Tumor Agnostic Testing)	7
Tumor-agnostic testing for patients with advanced solid tumors	7
Cancer-specific Criteria	9
Bladder Cancer (Urothelial Carcinoma, including the Upper Tract)	9
Breast Cancer	10
Cholangiocarcinoma (Biliary Tract Cancers)	12
Colorectal Cancer.....	13
Melanoma.....	14
Non-Small Cell Lung Cancer	16
Ovarian cancer (epithelial).....	17
Pancreatic Adenocarcinoma.....	18
Prostate Cancer	18
Thyroid Cancer	20
Unknown Primary Site Cancer	22
Somatic Testing of Hematologic Malignancies	23
Clinical Indications	23
General Requirements	23
Umbrella Criteria	23
Somatic Genomic Testing (Tumor Biomarker Testing).....	23
Cancer-specific Criteria	24
Acute Lymphocytic Leukemia	24
Acute Myelogenous Leukemia	25
Chronic Myeloid Leukemia and Myeloproliferative Disorders	25
Multiple Myeloma	27
Myelodysplastic Syndrome.....	27
References.....	28
Codes.....	35
History	42

Description and Application of the Guidelines

The Carelon Clinical Appropriateness Guidelines (hereinafter “the Carelon Clinical Appropriateness Guidelines” or the “Guidelines”) are designed to assist providers in making the most appropriate treatment decision for a specific clinical condition for an individual. As used by Carelon, the Guidelines establish objective and evidence-based criteria for medical necessity determinations where possible. In the process, multiple functions are accomplished:

- To establish criteria for when services are medically necessary (i.e., in general, shown to be effective in improving health outcomes and considered the most appropriate level of service)
- To assist the practitioner as an educational tool
- To encourage standardization of medical practice patterns
- To curtail the performance of inappropriate and/or duplicate services
- To advocate for patient safety concerns
- To enhance the quality of health care
- To promote the most efficient and cost-effective use of services

The Carelon guideline development process complies with applicable accreditation standards, including the requirement that the Guidelines be developed with involvement from appropriate providers with current clinical expertise relevant to the Guidelines under review and be based on the most up-to-date clinical principles and best practices. Relevant citations are included in the References section attached to each Guideline. Carelon reviews all of its Guidelines at least annually.

Carelon makes its Guidelines publicly available on its website twenty-four hours a day, seven days a week. Copies of the Carelon Clinical Appropriateness Guidelines are also available upon oral or written request. Although the Guidelines are publicly-available, Carelon considers the Guidelines to be important, proprietary information of Carelon, which cannot be sold, assigned, leased, licensed, reproduced or distributed without the written consent of Carelon.

Carelon applies objective and evidence-based criteria, and takes individual circumstances and the local delivery system into account when determining the medical appropriateness of health care services. The Carelon Guidelines are just guidelines for the provision of specialty health services. These criteria are designed to guide both providers and reviewers to the most appropriate services based on a patient’s unique circumstances. In all cases, clinical judgment consistent with the standards of good medical practice should be used when applying the Guidelines. Guideline determinations are made based on the information provided at the time of the request. It is expected that medical necessity decisions may change as new information is provided or based on unique aspects of the patient’s condition. The treating clinician has final authority and responsibility for treatment decisions regarding the care of the patient and for justifying and demonstrating the existence of medical necessity for the requested service. The Guidelines are not a substitute for the experience and judgment of a physician or other health care professionals. Any clinician seeking to apply or consult the Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient’s care or treatment.

The Guidelines do not address coverage, benefit or other plan specific issues. Applicable federal and state coverage mandates take precedence over these clinical guidelines. If requested by a health plan, Carelon will review requests based on health plan medical policy/guidelines in lieu of the Carelon Guidelines. Pharmaceuticals, radiotracers, or medical devices used in any of the diagnostic or therapeutic interventions listed in the Guidelines must be FDA approved or conditionally approved for the intended use. However, use of an FDA approved or conditionally approved product does not constitute medical necessity or guarantee reimbursement by the respective health plan.

The Guidelines may also be used by the health plan or by Carelon for purposes of provider education, or to review the medical necessity of services by any provider who has been notified of the need for medical necessity review, due to billing practices or claims that are not consistent with other providers in terms of frequency or some other manner.

General Clinical Guideline

Clinical Appropriateness Framework

Critical to any finding of clinical appropriateness under the guidelines for a specific diagnostic or therapeutic intervention are the following elements:

- Prior to any intervention, it is essential that the clinician confirm the diagnosis or establish its pretest likelihood based on a complete evaluation of the patient. This includes a history and physical examination and, where applicable, a review of relevant laboratory studies, diagnostic testing, and response to prior therapeutic intervention.
- The anticipated benefit of the recommended intervention should outweigh any potential harms that may result (net benefit).
- Current literature and/or standards of medical practice should support that the recommended intervention offers the greatest net benefit among competing alternatives.
- Based on the clinical evaluation, current literature, and standards of medical practice, there exists a reasonable likelihood that the intervention will change management and/or lead to an improved outcome for the patient.

If these elements are not established with respect to a given request, the determination of appropriateness will most likely require a peer-to-peer conversation to understand the individual and unique facts that would supersede the requirements set forth above. During the peer-to-peer conversation, factors such as patient acuity and setting of service may also be taken into account.

Simultaneous Ordering of Multiple Diagnostic or Therapeutic Interventions

Requests for multiple diagnostic or therapeutic interventions at the same time will often require a peer-to-peer conversation to understand the individual circumstances that support the medical necessity of performing all interventions simultaneously. This is based on the fact that appropriateness of additional intervention is often dependent on the outcome of the initial intervention.

Additionally, either of the following may apply:

- Current literature and/or standards of medical practice support that one of the requested diagnostic or therapeutic interventions is more appropriate in the clinical situation presented; or
- One of the diagnostic or therapeutic interventions requested is more likely to improve patient outcomes based on current literature and/or standards of medical practice.

Repeat Diagnostic Intervention

In general, repeated testing of the same anatomic location for the same indication should be limited to evaluation following an intervention, or when there is a change in clinical status such that additional testing is required to determine next steps in management. At times, it may be necessary to repeat a test using different techniques or protocols to clarify a finding or result of the original study.

Repeated testing for the same indication using the same or similar technology may be subject to additional review or require peer-to-peer conversation in the following scenarios:

- Repeated diagnostic testing at the same facility due to technical issues
- Repeated diagnostic testing requested at a different facility due to provider preference or quality concerns
- Repeated diagnostic testing of the same anatomic area based on persistent symptoms with no clinical change, treatment, or intervention since the previous study

- Repeated diagnostic testing of the same anatomic area by different providers for the same member over a short period of time

Repeat Therapeutic Intervention

In general, repeated therapeutic intervention in the same anatomic area is considered appropriate when the prior intervention proved effective or beneficial and the expected duration of relief has lapsed. A repeat intervention requested prior to the expected duration of relief is not appropriate unless it can be confirmed that the prior intervention was never administered.

Somatic Testing of Solid Tumors

Clinical Indications

General Requirements

Repeated testing of the same individual for the same indication should be limited to evaluation following an intervention, or when there is a change in clinical status such that additional testing is required to determine next steps in management. At times, it may be necessary to repeat a test using different techniques or protocols to clarify a finding or result of the original study.

Repeated testing for the same indication using the same or similar technology may be subject to additional review or require peer-to-peer conversation in the following scenarios:

- Repeated diagnostic testing of the same tumor site with no clinical change, treatment, or intervention since the previous study
- Repeated diagnostic testing of the same individual and the same tumor by different providers for the same member over a short period of time

Umbrella Criteria

Somatic Genomic Testing (Tumor Biomarker Testing)

Somatic genomic testing is considered **medically necessary** in individuals with cancer when **ANY** of the following criteria are met:

- The genomic testing has established analytical and clinical validity and is performed in an appropriately certified laboratory
- The genetic test has established clinical utility such that a positive or negative result will meaningfully impact the clinical management (predictive, diagnostic, prognostic, or therapeutic) of the individual and will likely result in improvement in net health outcomes (i.e., the health benefits of the interventions outweigh any medical or psychological harmful effects of the testing intervention)
 - When there are genomic biomarker-linked therapies approved by the U.S. Food and Drug Administration (FDA) for the individual's specific cancer scenario and such therapies are being considered in the near term
 - When considering a treatment for which there are specific genomic biomarker-based contraindications or exclusions related therapeutic decisions being considered in the near term
- Clinical decision making incorporates the known or predicted impact of a specific genomic alteration on protein expression or function and published clinical data on the efficacy of targeting that genomic alteration with a particular agent

Rationale

Nearly every malignancy will have somatic mutations that have been described, although most known mutations do not have clinical management implications. While various common conditions are covered by specific guideline criteria for somatic testing of tumors, it is not feasible to establish criteria for every clinical scenario in oncology and hematology. The general criteria for somatic testing (above) apply to malignancy when more specific criteria are not available.

Metastatic or Advanced Cancer (Tumor Agnostic Testing)

Tumor-agnostic testing for patients with advanced solid tumors

Multi-gene panel testing is considered **medically necessary** when **ALL** of the following are true:

- The individual has a metastatic or advanced solid tumor and adequate performance status for cancer treatment
- A genomic biomarker-linked therapy has been approved by the FDA for their cancer clinical scenario, or there are established genomic biomarker-based treatment contraindications or exclusions
- There are no existing indications for the planned therapy such that its use does not depend on the results of genetic testing (i.e., immune checkpoint inhibitor indications)
- There are no satisfactory tumor-specific standard therapies available
- Testing falls into **ANY** of the following categories:
 - Mismatch-repair (MMR) deficiency
 - MLH1, MSH2, MSH6, PMS2 or EPCAM genes by PCR or NGS testing
 - Microsatellite testing (MSI)
 - MMR by IHC
 - Tumor mutational burden (TMB) testing
 - NTRK fusion testing
 - BRAF V600E mutation testing

Rationale

Oncologists have traditionally chosen therapies and determined prognoses based on site of origin and histology. In select tumor types, oncologists began incorporating biomarkers, such as immunohistochemistry (IHC) for HER2 and estrogen/progesterone receptor status in breast cancer into their decision-making.¹ Today, genomic characterization is increasingly being used to guide treatment decisions, especially in patients with advanced disease. Large-scale sequencing studies such as those of The Cancer Genome Atlas and the International Cancer Genome Consortium have described the genomic landscape of 20-30 solid tumor types, identifying certain alterations as drivers. Subsequent studies have defined a consensus list of cancer driver genes and patterns of co-occurrence and mutual exclusivity of these alterations.² Studies of comprehensive NGS testing in patients with advanced cancer report a wide range of clinically actionable genomic alterations per patient, ranging from 40% to 94%. Furthermore, only 10%-25% of patients actually receive therapy informed by sequencing.³ The only randomized clinical trial to explore the clinical effects of delivering genomically directed therapy to patients undergoing NGS testing in the setting of advanced cancer found no improvement in progression-free survival for patients receiving molecularly matched therapy.⁴ Thus, rather than systematic NGS testing in all advanced cancer patients, the current standard of care involves somatic testing applied in various specific tumor scenarios where such testing is known to be important because of driver alterations that are effectively treated with targeted agents.

In addition to the specific tumor scenarios where NGS testing is indicated, the FDA has also approved the use of treatment regimens for tumor-agnostic indications in several specific scenarios where patients have progressed following previous treatment and have no satisfactory alternative treatments: pembrolizumab for patients with microsatellite instability (since 2017) or high tumor mutational burden (since 2020), larotrectinib (since 2018) or entrectinib (since 2019) for use in patients harboring tumors with NTRK fusions, and dabrafenib plus trametinib (since 2022) when the tumor harbors a BRAF V600E mutation.

Microsatellite instability (MSI) is the result of inactivation of the DNA mismatch repair (MMR) system and is characterized by a high frequency of frameshift mutations in microsatellite DNA. In a portion of tumors, MSI is caused by germline mutations in one of the MMR genes (MLH1, MSH2, MSH6, or PMS2), which results in hereditary Lynch syndrome. However, the majority (80%) of MSI cases are sporadic, often because of hypermethylation of the MLH1 gene promoter.⁵ The diagnosis of microsatellite instability (MSI) and mismatch repair deficiency (dMMR) status is made using two reference techniques: molecular biology (polymerase chain reaction) and immunohistochemistry, with 90% to 97% agreement between the two techniques; the sensitivity of next-generation sequencing-based algorithms may not be as robust as immunohistochemistry and polymerase chain reaction.⁶ The FDA did not specify which assay should be used to assess for mismatch repair or

microsatellite instability, and the College of American Pathologists in collaboration with several other organizations have guidelines pending. In the draft guidelines, for patients being considered for use of checkpoint inhibitors with cancer types other than colorectal, gastroesophageal, small bowel or endometrial cancer, the recommendation is that pathologists should test MMR genes for DNA mismatch repair, although the optimal approach is unknown. Overall, DNA mismatch repair is found in roughly 4% of all adult cancer.⁵

The FDA granted tissue-agnostic accelerated approval in June 2020 for the anti-PD-1 pembrolizumab in TMB ≥ 10 mutations/Mb solid tumors as determined by an FDA-approved test, for patients that have progressed following prior treatment and who have no satisfactory alternative treatment options. TMB testing is based on the underlying assumption that increasing the numbers of mutant proteins will create antigenic peptides allowing for enhanced immunogenicity.⁷ The conceptual definition of TMB is total number of mutations present in a tumor specimen. The actual definition of the type of genetic alterations considered for TMB has varied according to different methodologies.⁸ The FoundationOneCDx assay (Foundation Medicine, Inc., CPT 0037U) was the FDA approved a companion diagnostic. The MSK-IMPACT test was cleared through the FDA 510(k) process in 2017 and in the following years other NGS panels have been approved this way too. The Foundation Medicine TMB assay uses a tissue-based specimen and examines a genomic region of approximately 1.1 Mb. For TMB estimation this test includes synonymous and non-synonymous mutations and short indels, while oncogenic drivers are excluded. In addition, germline alterations are excluded based on validated bioinformatics algorithms. There are at least many other TMB tests in the market, including tests from Memorial Sloan Kettering Cancer Center (MSK-IMPACT), Tempus, Qiagen, Neogenomics, Illumina, Guardant Health, Caris, Thermo-Fisher Scientific, and others. Each lab tests a different number of genes, covers a different total region of the genome, and some use only nonsynonymous mutations. Each test uses its own bioinformatics algorithm and the type of specimens used also varies. The equivalence of 10 mut/Mb in FoundationOne CDx cannot be easily determined in other TMB panels. For example, the cutoff of high TMB for the MSK-IMPACT assay was defined at 13.8 mut/Mb based on their cohort.⁹ Further alignment in the numerical cutoff of TMB across panels requires calibration tools and reference datasets or materials. The variation between TMB estimates can confuse clinicians and may hinder clinical decision making.¹⁰

The decision for the FDA to provide accelerated approval drew mixed reactions from the oncology community. On one hand, this was seen as a positive development because of the strong biological rationale for high TMB as a biomarker for immunotherapy sensitivity, and this was felt to address an unmet need for patients with rare cancer types to improve their access to immunotherapy.¹¹ Critics noted that 10 mut/Mb is an arbitrary cut off that does not separate responders from non-responders and it has not been shown to be associated with improved overall survival or quality of life. Additionally, there were concerns that there is insufficient evidence across tumor types for which pembrolizumab was not already approved and insufficient evidence to know if pembrolizumab is more cost effective than alternatives.¹² PD-L1 expression and TMB are not significantly correlated within most cancer subtypes, and TMB may not always correlate with ICI responsiveness.⁸ Data examined from over 10,000 patients included in the Cancer Genome Atlas with TMB measured by the FoundationOne CDx assay failed to support the use of TMB-H as a biomarker for immune checkpoint blockade treatment in all tumor types, including at the FDA-approved threshold of 10 mut/Mb.⁷ There are still many challenges for the further development of TMB as biomarker with clinical utility. Prospective randomized trials are required to establish the roles of TMB and other ICI biomarkers in a variety of clinical settings. For instance, the predictive value of TMB for combinations of immunotherapies with targeted agents or chemotherapy is not established.

Members of the neurotrophic receptor tyrosine kinase (NTRK) fusion oncogene family, NTRK1/NTRK2/NTRK3, are most prevalent in rare adult cancer types and in several pediatric cancers, although they can occur in a very small proportion of commonly occurring cancer types in adults, including NSCLCs, CRCs, head and neck cancers, thyroid cancers (especially in those with a history of radiation exposure, gliomas, inflammatory myofibroblastic tumors and some other sarcomas, and melanocytic tumors.^{5, 13} Evidence of how rare NTRK fusions are comes from an examination of tissue samples from 11,502 patients where 53 gene fusions were analyzed and sequencing of 592 genes was done along with an IHC evaluation of TrkA/B/C. This review found only thirty-one cases (0.27% of the entire cohort) with NTRK fusions.¹⁴ Nevertheless, when NTRK fusions are found, larotrectinib and entrectinib have shown to have significant and durable activity against locally advanced and metastatic solid tumors with NTRK fusions.^{14, 15} Notably, this activity was seen regardless of tumor site of origin, histologic classification, or NTRK fusion type.

Immunohistochemistry (IHC) is commonly used for NTRK testing and is practical to implement in most laboratories. IHC has variable specificity according to tumour type. While the antibody appears to have 100% specificity in carcinomas of the colon, lung, thyroid, pancreas and biliary tract, decreased specificity is seen in breast and salivary gland carcinomas, as cytoplasmic staining can occasionally be seen. Specificity is lower in sarcomas, particularly those with neural or smooth muscle differentiation.¹³ Fluorescence in situ hybridization (FISH) can detect large structural variants at the DNA level. A commercial break-apart probe is available for the ETV6 gene. NGS methods can also be used. NGS testing can be particularly useful for monitoring patients with NTRK fusions for development of resistance mutations. There are some limitations of using NGS testing, however. For example, the sensitivity of DNA-based NGS suffers if fusion breakpoints involve long intronic regions. Also, when novel structural variants are detected, it can be difficult to determine whether such an event results in a functional expressed fusion. Other drawbacks include turnaround time (typically several weeks, and that more material is required for testing. Selection of the appropriate assay for NTRK fusion detection therefore depends on tumor type and genes involved, as

well as consideration of other factors such as available material, accessibility of various clinical assays, and whether comprehensive genomic testing is needed concurrently.¹⁶

The European Society for Medical Oncology (ESMO) Translational Research and Precision Medicine Working Group launched a collaborative project to propose a classification system for molecular aberrations based on the evidence available supporting their value as clinical targets and established from this work the ESMO Scale for Clinical Actionability of molecular Targets (ESCAT).¹⁷ The ESCAT acknowledges that for the qualification of (ultra-) rare molecular aberrations it might not be feasible to obtain data from prospective randomized clinical trials (RCTs). Tier I, consisting of targets suitable for routine clinical use, therefore has two subclasses with targets for which alternative evidence is available: tier I-B includes "targets are supported by data from prospective, non-randomized clinical trials that, while unable to provide evidence for survival improvement, have demonstrated clinically meaningful benefit as defined by the ESMO Magnitude of Clinical Benefit Scale"¹⁷ and IC includes targets for which "clinical trials in multiple tumor types, or basket clinical trials, have demonstrated a clinically meaningful benefit for the target-drug pair with similar magnitude of benefit across the different tumor types."¹⁸ Based on ESCAT, NTRK fusions, high TMB status, and microsatellite instability (MSI) are typically ranked IC in this rubric for most solid tumors.

Finally, the FDA granted accelerated approval in June 2022 to dabrafenib in combination with trametinib for the treatment of adult and pediatric patients ≥ 6 years of age with unresectable or metastatic solid tumors with BRAF V600E mutation who have progressed following prior treatment and have no satisfactory alternative treatment options. Dabrafenib in combination with trametinib is not indicated for patients with colorectal cancer because of known intrinsic resistance to BRAF inhibition. Mutations in BRAF occur in many tumor types and contribute to the dysregulation of processes such as cell proliferation and differentiation. Acquired resistance is common among patients receiving BRAF inhibitor monotherapies. Efforts to overcome this in BRAF V600 mutation-positive melanoma, NSCLC, and ATC have tested combined MEK and BRAF inhibition.¹⁹ The FDA approval was based on safety and efficacy evaluation in 131 adult patients from open-label, multiple cohort trials BRF117019 and NCI-MATCH, 36 pediatric patients from CTMT212X2101, and supported by results in COMBI-d, COMBI-v, and BRF113928. In addition, there have been data published from the phase 2, open-label, single-arm, multicentre, Rare Oncology Agnostic Research (ROAR) basket trial in patients with BRAFV600E-mutated rare cancers with promising findings found in thyroid cancer²⁰, malignant gliomas²¹, and biliary tract cancer.²²

Cancer-specific Criteria

Bladder Cancer (Urothelial Carcinoma, including the Upper Tract)

Tissue-based somatic tumor testing for FGFR mutations is considered **medically necessary** for individuals with urothelial tumors of the bladder or upper urinary tract when **ALL** of the following criteria are met:

- The individual has biopsy-proven urothelial malignancy
- The urothelial malignancy is locally advanced or metastatic
- An FDA-approved RT-PCR FGFR testing kit is used
- The individual is a potential candidate for targeted therapy prescribed on the basis of the FGFR test results
- The individual has not had prior FGFR testing in the metastatic setting

Tissue-based somatic tumor testing for microsatellite instability (MSI testing) is considered **medically necessary** for individuals with muscle-invasive urothelial tumors of the upper urinary tract.

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Bladder cancers are biologically diverse and can be separated into "molecular subtypes," based on expression profiling.^{23, 24} In these molecular classification systems, over 90% of muscle-invasive bladder cancers classify as either luminal or basal-squamous subtypes. These subtypes associate with clinical behavior, histology, and molecular alterations, though their clinical utility has not been demonstrated at present and use in bladder cancer is not recommended.²⁵

Oncogenic alteration of FGFR3 is present in approximately 15% of muscle-invasive bladder cancers. The luminal subtype of bladder cancer is enriched in FGFR3 mutations and FGFR3 overexpression.²⁵ Also, FGFR mutations are more frequent in the upper tract ($\approx 30\%$) than the bladder ($\approx 14\%$).²⁶ A phase 2 study (BLC2001) in 99 patients with locally advanced and metastatic disease who did not respond to prior therapy found a 40% objective response rate with oral erdafitinib, a pan-FGFR inhibitor.²⁷

The FDA granted accelerated approval of this drug in patients with advanced or metastatic urothelial carcinoma, with relevant FGFR alterations, whose disease has progressed during or following treatment with platinum-based chemotherapy, including in the adjuvant and neoadjuvant settings. Testing may be performed using the FDA-approved companion diagnostic (a specific RT-PCR kit), or other methods, including next-generation sequencing.²⁵ The International Society of Urologic Pathology (ISUP) working group does not recommend testing all patients with advanced-stage disease at the time of diagnosis. Instead, the IUSP recommends FGFR testing be performed more selectively, on patients with advanced disease who have progressed following platinum-based therapy, or who have another indication to perform testing, based on the judgment of treating physicians.

In a series of patients with advanced upper tract urothelial cancer (479 patients) and bladder urothelial cancer (1984 patients), comprehensive genomic profiling revealed 0.7% of the bladder cancers and 3.1% of the upper tract cancers were MSI-H. FGFR3 activating mutations or fusions were found in 18% of the bladder tumors and 25.5% of the upper tract tumors.²⁸

Breast Cancer

Localized breast cancer

Gene expression profiling is considered **medically necessary** for individuals with localized breast cancer using Oncotype DX, MammaPrint, EndoPredict, Prosigna Breast Cancer Prognostic Gene Signature Assay, or the Breast Cancer Index when **ALL** of the following criteria are met:

- Surgery has been performed and a full pathological evaluation of the specimen has been completed
- Histology is ductal, lobular, mixed, or metaplastic
- Receptor status is estrogen receptor positive (ER+), progesterone receptor positive (PR+), or both; **AND** HER2-negative
- Lymph node status is node-negative (pN0) or axillary lymph node micro-metastasis (pN1mi) less than or equal to 2 mm
- Tumor features include **ANY** of the following:
 - Tumor size greater than 1.0 cm and less than or equal to 5.0 cm
 - Tumor size 0.6–1.0 cm and moderate or poorly differentiated
 - Tumor size 0.6–1.0 cm and well-differentiated with **ANY** of the following:
 - angiolymphatic invasion
 - high nuclear grade
 - high histologic grade
- Chemotherapy is being considered by the individual and their provider
- No other breast cancer gene expression profiling assay has been conducted for this tumor (this includes testing on any metastatic foci or on other sites when the tumor is multifocal)

Gene expression profiling with the Oncotype DX or MammaPrint is considered **medically necessary** for postmenopausal individuals with 1 to 3 positive axillary lymph nodes (pN1a, pN1b or pN1c) when **ALL** of the following criteria are met:

- Surgery has been performed and a full pathological evaluation of the specimen has been completed
- Histology is ductal, lobular, mixed, or metaplastic
- Receptor status is estrogen receptor positive (ER+), progesterone receptor positive (PR+), or both; **AND** HER2-negative
- Chemotherapy is being considered by the individual and their provider
- No other breast cancer gene expression profiling assay has been conducted for this tumor (including testing on any metastatic foci or on other sites when the tumor is multifocal)

Metastatic breast cancer

Testing for somatic pathogenic variants of PIK3CA is considered **medically necessary** when **ALL** of the following criteria are met:

- The individual has ER-positive and HER2-negative metastatic breast cancer
- The individual is a candidate for alpelisib or another FDA-approved PIK3CA-targeted agent
- The individual has not had prior testing for PIK3CA in the metastatic setting

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Adjuvant therapy for breast cancer

Breast tumors are routinely pathologically evaluated by immunohistochemical staining for the presence of ER, PR, and HER2 overexpression. This information is used to make treatment decisions about hormonal and HER2-targeted therapy. Characterizing the tumor's gene expression profile allows for risk stratification for recurrence. Multigene expression assays are currently used in the care of more than one-third of women with breast cancer in the United States.²⁹ These genetic profiling tests are used to assess the benefit of adjuvant chemotherapy for early-stage breast cancer and ASCO has published a guideline about the use of biomarkers for adjuvant endocrine and chemotherapy use since 2016. The most commonly used test for this purpose is Oncotype DX, a 21-gene assay. The sentinel trial establishing its clinical utility for use in women with lymph node negative early breast cancer, the TailorX trial, enrolled 10,273 women and was published in 2018³⁰ with earlier data about its prospective validation published in 2015³¹ and subsequent data showing how other clinical data added to this test further refined its use.³⁰ A separate large clinical trial, the RXPonder trial which enrolled 5,083 women addressed the clinical utility of the Oncotype DX test in women with early stage, lymph node positive breast cancer.³² Moreover, the clinical utility of another multigene assay, the 70-gene signature MammaPrint, was evaluated in a prospective randomized trial (MINDACT), that was published in 2016. This trial included 6,693 women with node-negative or 1-3 node-positive, early-stage breast cancer.³³ Patients were eligible irrespective of breast cancer subtype, but the majority had ER-positive tumors. Subsequent analyses of MINDACT further refinement of its use based on age.³⁴ The most recent update of the ASCO guideline in 2022³⁵ categorizes these clinical scenarios based on the lymph node status (node-negative vs. 1-3 positive nodes vs. ≥ 4 nodes), age and menopausal status (premenopausal or age ≤ 50 years vs. other), and HER2 receptor status. There is insufficient evidence in the realms of HER2 positive disease and for patients with ≥ 4 nodes. There is high quality evidence and a strong recommendation for use of Oncotype DX in node-negative women (regardless of age or menopausal status) and in those with 1-3 positive lymph nodes. For MammaPrint, the ASCO guideline indicates intermediate levels of evidence and strong recommendation for node-negative women who are postmenopausal, age ≥ 50 years, or those with 1-3 positive lymph nodes. For these same categories noted above for MammaPrint, a lower level recommendation (moderate) was given to several additional tests including EndoPredict, Prosigna, Breast Cancer Index, and use of non-genetic tests like IHC4 (which combines ER, PR, HER2 and Ki67 into one score) or the Ki-67 test. There is no current role for emerging biomarkers in this setting such as PD-L1 testing, evaluation of circulating tumor cells, or measurement of tumor-infiltrating lymphocytes.

The ASCO guideline update on biomarker use in early-stage breast cancer also that extended adjuvant endocrine therapy (beyond 5 years) has demonstrated improved outcomes albeit with modest absolute benefit and added toxicity and tolerability challenges. In clinical practice, the decision about extended adjuvant therapy is a shared decision negotiated between the patient and physician at the time of the clinical decision (typically within 3 months or so of the time that the 5-year course of adjuvant hormonal therapy is completed). This shared decision-making is highly personal and it may take into account the individual patient's risk of recurrence with or without further adjuvant therapy, fear of recurrence, the physical, psychological and financial tolerability of the adjuvant hormonal therapy, competing risks to health, and sometimes other factors. This is a clinical scenario where prospective data evaluating the clinical utility of the incremental information based on genomics is particularly crucial. Unfortunately, there is no prospective data evaluating whether or not the use of the a multigene classifier like the Breast Cancer Index, based on its ability to refine the risk of recurrence estimates above and beyond what is already known, adds clinical utility to that shared decision in terms of better health outcomes, less distress or decisional regret, less anxiety, or other patient-centered outcomes.

In the ASCO guideline update on biomarker use in breast cancer, on a scale ranging from "should use" to "may offer" to "insufficient evidence" to "should not use", the Breast Cancer Index landed with a "may use" statement based on intermediate quality evidence from the Ideal study and two smaller prospective-retrospective studies.^{36, 37} These studies each showed evidence of prognostic associations but no prospective data addressing clinical utility such as seen with trials like TailorX, RXPonder, or MindACT in the different clinical setting of initial adjuvant therapy decision-making.

Metastatic breast cancer

The European Society for Medical Oncology (ESMO) Translational Research and Precision Medicine Working Group launched a collaborative project to propose a classification system for molecular aberrations based on the evidence available supporting their value as clinical targets and established from this work the ESMO Scale for Clinical Actionability of molecular Targets (ESCAT).¹⁷ The ESCAT acknowledges that for the qualification of (ultra-) rare molecular aberrations it might not be feasible to obtain data from prospective randomized clinical trials (RCTs). Tier I, consisting of targets suitable for routine clinical use, therefore has two subclasses with targets for which alternative evidence is available: tier I-B includes ‘targets are supported by data from prospective, non-randomized clinical trials that, while unable to provide evidence for survival improvement, have demonstrated clinically meaningful benefit as defined by the ESMO Magnitude of Clinical Benefit Scale and IC includes targets for which ‘clinical trials in multiple tumor types, or basket clinical trials, have demonstrated a clinically meaningful benefit for the target-drug pair with similar magnitude of benefit across the different tumor types.’¹⁸ Based on the current evidence, the ESMO Precision Medicine working group does not recommend routine use of large panel NGS testing on tumor samples for patients with metastatic breast cancer.³⁸ HER2 amplification, germline BRCA1/2 mutations, and PIK3CA mutations were classified in tier of evidence IA based on large randomized trials showing antitumor activity of targeted therapies in patients presenting with these alterations. Neurotrophic receptor tyrosine kinase (NTRK) fusions and microsatellite instability (MSI) were ranked IC.³⁹

Genotype-driven targeted therapy using alpelisib has received regulatory approval and become part of routine clinical practice for PIK3CA-mutated ER-positive/HER2-negative metastatic breast cancer based on the Solar-1 trial data.⁴⁰ Approximately 40% of patients with ER-positive/HER2-negative disease have activating PIK3CA mutations. In Solar-1, the presence or absence of any PIK3CA mutation by means of polymerase-chain-reaction analysis of mutation hot spots in the C2, helical, and kinase domains of PI3K (corresponding to exons 7, 9, and 20, respectively) with the use of a tumor-tissue sample. In addition, for patients harboring germline BRCA1 or BRCA2 mutations, the use of olaparib (a PARP inhibitor) is now a standard treatment option based on data from the OlympiAD trial.⁴¹ Based on these data, the presence of metastatic breast cancer is an indication for germline BRCA testing, although many patients have already undergone such testing by the time they develop metastatic disease since de novo metastatic breast cancer represents less than 10% of metastatic breast cancer and the indications for BRCA testing are broad.

Most therapies targeting kinase fusions are either still in clinical development or approved in tumor types other than breast cancer except for the NTRK inhibitors which received tumor-agnostic regulatory approval and would be an option for 4 out of 4854 patients (0.08%) of patients with breast cancer (Hilbers). While somatic ESR1 mutations are a well-established mechanism of acquired resistance to aromatase inhibitors, clinical trials have demonstrated that these aromatase inhibitors combined with CDK 4/6 agents remain active in the presence of ESR1 mutations.⁴² Other biomarkers such as AKT1, FGFR1/FGFR2, PTEN, NFI, and tumor signatures (APOBEC) are still under investigation in clinical trials.

Cholangiocarcinoma (Biliary Tract Cancers)

Tissue-based NGS panel testing of tumors in individuals with cholangiocarcinoma is considered **medically necessary** when **ALL** of the following criteria are met:

- The individual has biopsy-proven cholangiocarcinoma
- The cholangiocarcinoma is locally advanced or metastatic
- The panel testing contains, at minimum, testing of pathogenic variants in the following genes: IDH1, FGFR, and BRAF
- The individual is a potential candidate for targeted therapy prescribed on the basis of the panel test results
- The individual has not had prior NGS testing in the metastatic setting

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Biliary tract cancer refers to a spectrum of invasive adenocarcinomas, including cholangiocarcinoma (cancers arising in the intrahepatic, perihilar, or distal biliary tree), and gallbladder carcinoma. Identification of distinct patient subgroups with driver mutations amenable to targeted therapy have been identified, which are generally mutually exclusive from one another and often associated with the anatomical subsite of the tumor.⁴³ The most common molecular alterations in biliary tract malignancy are found in IDH1 (mutations found in 13% of intrahepatic cholangiocarcinoma cases), the FGFR pathway, particularly FGFR2

mutations (20% of intrahepatic cholangiocarcinoma), HER2 mutations (found in up to 20% of gallbladder and extra-hepatic cholangiocarcinoma), BRAF V600E mutations (found in 5% of intrahepatic cholangiocarcinoma). To identify these potentially targetable alterations, performing molecular profiling for patients diagnosed with cholangiocarcinoma (especially intrahepatic cholangiocarcinoma) is now considered standard of care.⁴⁴ Like other solid tumors, mutations in NTRK or MMR deficiency may also be found rarely.

A majority of patients are diagnosed with advanced disease, when chemotherapy with cisplatin and gemcitabine followed by second-line chemotherapy is the cornerstone of treatment for most patients in the absence of targetable alterations. For patients with advanced IDH-1-mutant cholangiocarcinoma refractory to one or two lines of systemic therapy, the IDH-1 inhibitor ivosidenib has shown activity in the ClarIDHy phase III clinical trial when compared with placebo.⁴⁵ For patients with FGFR2 fusions or other rearrangements, phase II single-arm registrational trials of FGFR inhibitors in the previously treated, unresectable locally advanced or metastatic cholangiocarcinoma population show an overall response rate in the 23%-42% range and a median progression-free survival of 7 to 9 months and FGFR inhibitors such as pemigatinib and infigratinib can be used in this setting. Likewise, there is activity in treating patients with BRAF V600E mutations with dabrafenib plus trametinib as well as use of entrectinib to treat those with NTRK inhibitors. HER2-directed therapies have some activity, albeit less convincing, in chemo-refractory patients.⁴⁴

Colorectal Cancer

Localized colorectal cancer

Tissue-based somatic tumor testing is considered **medically necessary** for individuals with localized (stage II-III) colorectal cancer when **ALL** of the following criteria are met:

- The individual has biopsy-proven adenocarcinoma of the colon or rectum
- Includes any or all of the following, with no prior testing
 - MSI testing
 - BRAF V600E
 - MMR testing by IHC

See Hereditary Cancer Testing guideline for further details regarding indications for germline MMR testing.

Metastatic colorectal cancer

Tissue-based somatic tumor testing is considered **medically necessary** for individuals with metastatic colorectal cancer and may be performed on the primary tumor or a metastatic site when **ALL** of the following criteria are met:

- The individual has biopsy-proven adenocarcinoma of the colon or rectum
- Assessment includes the following:
 - MSI testing and/or MMR testing (using IHC)
 - Extended RAS testing (KRAS and NRAS)
 - BRAF V600E
 - HER2 testing
- There has been no prior testing

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Among people diagnosed with colon cancer, 20% have metastatic CRC, and 40% present with recurrence after previously treated localized disease.⁴⁶ Among those with localized CRC, approximately 15% are deficient in one or more DNA mismatch repair (MMR) proteins, with one-quarter of these resulting from Lynch syndrome.⁴⁷ About 90 to 95 percent of colorectal cancers in patients with Lynch syndrome show evidence of microsatellite instability (MSI).⁴⁸ In patients with early-stage

(especially stage II) CRC, MMR status provides important prognostic and predictive information, with MMR-deficiency being associated with both a good prognosis and a lack of efficacy from fluorouracil treatment.⁴⁹

Five percent of metastatic CRC tumors are MMR-D or MSI-H (either due to Lynch syndrome or sporadic mutations) and may respond to immunotherapy.⁴⁶ All CRC (localized and metastatic) should be tested for either mismatch repair deficiency (MMR-D) via immunohistochemistry or microsatellite instability-high (MSI-H) via polymerase chain reaction to screen for Lynch syndrome and/or to inform therapeutic options. In the metastatic disease setting, in addition to testing for microsatellite instability and MMR-deficiency, extended RAS testing has been the standard of care since the ASCO Provisional Clinical Opinion in 2015 based on evidence indicating that RAS mutations in exons 2 (codons 12 and 13), 3 (codons 59 and 61), and 4 (codons 117 and 146) of both KRAS and NRAS are predictive of resistance to treatment with monoclonal antibodies (such as cetuximab and panitumumab) targeting the epidermal growth factor receptor (EGFR).⁵⁰ These anti-EGFR antibody treatments should only be considered for treatment of patients with mCRC who are identified as having tumors with no mutations detected after such extended RAS mutation analysis.

BRAF V600E mutation is also routinely tested in metastatic colorectal cancer. This mutation leads to constitutive BRAF kinase activity and sustained MAPK pathway signaling in colon cancer. Other activating mutations in BRAF in CRC are very rare. The prevalence of this mutation is about 5%–10% in mCRC. BRAF mutation tumors have been associated with female gender, advanced age, proximal colon tumor location, T4 tumors, and poor response to standard chemotherapy.^{51, 52} In the BEACON study, BRAF and MEK directed therapy with encorafenib plus cetuximab improved overall survival and overall response rates in the metastatic setting after one or two lines of prior therapy compared with standard palliative chemotherapy.⁵³ Another biomarker under exploration is HER2 testing. The frequency of HER2 overexpression among patients with CRC is known to be around 5%. Thus far, phase II trials evaluating HER2 targeted therapy have produced mixed results, but such therapy is active when RAS and BRAF are wild-type and several clinical trials are ongoing.⁵⁴ Finally, NTRK gene fusions are extremely rare, occurring in approximately 0.35% of colorectal cancer and are more frequently found among patients with MMR-D tumors.⁵⁵

Melanoma

Diagnostic and prognostic testing in melanoma

Gene expression profiling of suspected or established cutaneous or uveal melanoma for diagnosis or prognostication is considered **not medically necessary**.

Somatic tumor testing in advanced melanoma

Tissue-based somatic tumor testing for **BRAF V600E** pathogenic variant by validated IHC, PCR, or NGS methods for individuals with resectable or unresectable stage III or stage IV melanoma or high-risk stage IIC melanoma is considered **medically necessary** when **BOTH** of the following criteria are met:

- The individual has biopsy-proven malignant melanoma
- Prior testing has not been performed

Tissue-based somatic tumor testing for individuals with resectable or unresectable stage III or stage IV melanoma or high-risk stage IIC melanoma that is **BRAF V600E wild type** is considered **medically necessary** when **ALL** of the following criteria are met:

- The individual has biopsy-proven malignant melanoma
- Prior testing has not been performed
- Testing includes **ANY** or **ALL** of the following:
 - KIT mutation testing
 - NRAS mutation testing
 - Additional BRAF mutation testing

Testing of individuals with **metastatic uveal melanoma** for **HLA-A*0201** using is considered **medically necessary** when **ALL** of the following criteria are met:

- The individual has biopsy-proven uveal melanoma and evidence of metastatic disease
- Prior testing for HLA-A*0201 has not been performed
- The individual is a candidate for treatment with tebentafusp

**Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.*

Rationale

Diagnosis of indeterminate melanocytic lesions

Light microscopic evaluation by a trained pathologist can provide an accurate diagnosis for the great majority of melanocytic lesions. There is, however, a small subset of melanocytic lesions that eludes appropriate classification by conventional light microscopy alone, preventing accurate prediction of clinical behavior and recommendations for appropriate treatment.⁵⁶ Ancillary tests such as comparative genomic hybridization (CGH) and fluorescence in situ hybridization (FISH), have been developed to help guide the diagnosis of ambiguous melanocytic proliferations. Although sometimes used, the correlation between various patterns of chromosomal abnormalities and outcomes are yet unclear.⁵⁶ Commercially available gene expression classifiers have also been developed and clinically validated. For example, a 23-gene classifier (MyPath Melanoma) was shown to have up to 92% sensitivity and 93% specificity at distinguishing benign from malignant lesions.⁵⁷⁻⁵⁹ The clinical utility related to use of this kind of testing has not been demonstrated. Prospective studies controlling the risk of bias and including clinically meaningful outcomes are needed to better understand the role of techniques used to clarify the diagnosis of the target subset of difficult to diagnose melanocytic lesions.

Prognostic classification of localized cutaneous melanoma

There are numerous clinicopathologic features that are accurately associated with melanoma outcome for localized disease (eg, Breslow thickness, ulceration, and tumor-infiltrating lymphocytes). Currently, prognostication based on pathologic features is quite robust.⁶⁰ There is interest, however, in using gene expression profiling (GEP) to categorize localized (stage I and II) cutaneous melanoma according to risk of metastasis and to better guide clinical decisions such as when to consider sentinel lymph node biopsy or adapt the surveillance intensity. Gene expression profiling associates the expression levels of many genes with an outcome, such as melanoma recurrence. Current GEP tests largely assign the patient's tumor to a specific prognostic class (eg, class 1 vs class 2, or low risk vs high risk) rather than calculating a specific calculated survival.⁶⁰ A published meta-analysis of the 31-gene GEP test focused on three selected studies plus data from a novel cohort of 211 patients showed that recurrence-free and distant metastasis-free survival rates were 91.4% and 94.1% for Class 1A patients and 43.6% and 55.5% for Class 2B patients.⁶¹ The methodology of this meta-analysis was criticized by academic dermatologists due to a lack of a pre-specified protocol, lack of adjustment for confounders and missing data, incomplete risk of bias assessment, publication bias, and financial conflicts of interest among the authors who were also meta-analyzing studies they also co-authored.⁶² Overall, the clinical utility of GEP classifiers in this setting remains uncertain. These GEP tests also need to be weighed against (or integrated with) the best available phenotypic models and other simple and widely available outcome algorithms.⁶⁰ For example, numerous studies have identified risk factors that may be predictive of positive SLN in thin melanoma, such as ulceration, high mitotic rate, and younger patient age. These risk factors are easily determined and inexpensive, and are now referred to as adverse features in guidelines for use in clinical staging and workup.⁶³

Prognostic classification of uveal melanoma

Uveal melanoma is a rare intraocular cancer that carries a 30%–50% risk of metastasis within 5 years of diagnosis. Metastases are observed most commonly in the liver and less frequently in the lungs and other organs. Post-treatment surveillance is based on individual judgement and consensus-based guidelines typically involves imaging to evaluate signs and symptoms as indicated, with consideration of blood testing of liver function and liver and chest, abdominal, and pelvic imaging every 6 to 12 months for 10 years. Patients considered high risk may be imaged more frequently, such as every 3 to 6 months over the first 5 years.

The risk of metastases from a uveal melanoma has traditionally been predicted based on the study of the morphologic and pathologic features of the tumor. Tumor thickness, diameter, location, presence of extraocular extension, and histopathology have been consistently shown to independently predict the likelihood of developing metastatic disease.⁶⁴ Cytogenetic investigations show that monosomy 3 is the most frequent karyotypic abnormality and is present in approximately 50% to 60% of patients. Monosomy of chromosome 3 and additional copies of 8q have been correlated with reduced survival.⁶⁵ A commercially available 15-gene expression profile (DecisionDX-UM). Registry data involving 89 patients enrolled from 4 institutions indicate that 39 of 49 lower risk patients were ultimately managed with lower intensity management.⁶⁶ Another uveal melanoma genetic test from Impact Genetics Inc. evaluates tumor specimens for complete or partial loss, duplications, or isodisomy of Chromosomes 1p, 3, 6, and 8 with genetic sequencing of the GNAQ and GNA11 genes is offered for select specimens.⁶⁴ Importantly, no survival benefit from the early detection of asymptomatic disease in uveal melanoma has been documented. In view of the lack of evidence for the benefit of any specific surveillance strategy, particularly for late relapse, recommendations for these patients and their providers remain uncertain.⁶⁵ Factors that are sometimes considered in decisions about surveillance include the patient's enhanced emotional well-being, the potential to identify oligometastatic disease amenable to surgery or other local therapies, the hope for decreased morbidity from advanced disease, and identification of eligibility for clinical trials that assess novel therapies for advanced uveal melanoma.⁶⁷ Multiple studies have shown that certain morphologic characteristics of melanoma tumors (size and location) independently predict metastasis,

beyond the data provided by either chromosomal analysis or GEP. Clinicians must integrate the entire clinical picture, including tumor size, location, and tumor growth velocity when interpreting the results of genetic analysis of an individual uveal melanoma tumor.⁶⁴ Ultimately, the selection of surveillance modality used varies according to local expertise and patient preferences.

Somatic tumor testing for resectable or unresectable stage III or stage IV melanoma

The majority of melanomas have mutations in genes associated with the MAPK pathway such as BRAF, NRAS, or NF1. Activation of the BRAF kinase via mutation is the most common, occurring in 40%–60% of cases, with mutated NRAS, occurring in 15%–30% of cases.⁶⁸ More than 90% of the BRAF mutations are V600E, with V600K mutations accounting for most of the remainder.⁶⁸ Currently, BRAF mutation status is the only biomarker that predicts a therapeutic response in advanced melanoma and testing for BRAF V600E mutation is the current standard of care in the setting of resectable or unresectable stage III or stage IV melanoma and also highly recommended for resected, high risk stage IIC disease.⁶⁹

The main cutaneous melanoma subtypes are associated with different mutational landscapes: frequently mutated genes aside from BRAF include CDKN2A, NRAS and TP53 in cutaneous melanoma, NRAS, NF1 and KIT in acral melanoma, and SF3B1 in mucosal melanoma.⁶⁹ In metastatic uveal melanoma, Testing for HLA-A*0201 by a specific assay used to identify patients who may be treated with tebentafusp (a bispecific protein directed to target glycoprotein 100 positive cells), which improved overall survival in a phase 3 trial.⁷⁰

NRAS mutations are present in 15% of melanomas and correlate with poor prognosis.⁷¹ BRAF-mutant and NRAS-mutant melanomas are sensitive to MEK inhibition, and the MEK inhibitor binimetinib has shown clinical activity in a controlled trial in the subset of patients with NRAS -mutant metastatic melanoma.⁷² Moreover, c-KIT mutations are also common and found in about 15%–23% of mucosal melanoma, 10%–15% of acral lentiginous melanoma (ALM), and 6%–28% of melanoma arising in chronically sun-damaged skin.⁷³ A review of 19 single arm studies of use of c-KIT inhibitors in unselected patients with KIT mutations or amplifications showed a response rate of 14% for mucosal melanoma and 22% for acral lentiginous melanoma, with activity in particular for mutations in exons 11 and 13 and minimal activity in KIT amplified tumors.⁷³ Overall, for patients in whom the melanoma is BRAF wild-type at the V600 locus, testing for less common BRAF mutations and for NRAS and c-KIT mutations is considered an ESMO evidence level IIC recommendation (suggesting that it is optional as there is insufficient evidence for efficacy).⁶⁹

Non-Small Cell Lung Cancer

Localized (stage IB-IIIa) NSCLC

Tissue-based somatic testing is considered **medically necessary** to identify EFGR pathogenic variant in individuals with localized NSCLC when **ALL** of the following criteria are met:

- Biopsy-proven, stage IB-IIIa NSCLC with **ANY** of the following characteristics:
 - An adenocarcinoma component on histology
 - Non-squamous, non-small cell histology
 - Any non-small cell histology when **EITHER** of the following clinical features are present:
 - Age 50 years or younger
 - Light or absent tobacco exposure
- Test results will determine candidacy for treatment with osimertinib

Metastatic NSCLC

Tissue-based NGS panel testing is considered **medically necessary** to identify pathogenic variants in individuals with stage IIIB, IIIC, or metastatic NSCLC when **ALL** of the following criteria are met:

- Biopsy-proven NSCLC with **ANY** the following characteristics:
 - An adenocarcinoma component on histology
 - Non-squamous, non-small cell histology
 - Any non-small cell histology when **EITHER** of the following clinical features are present:
 - Age 50 years or younger

- Light or absent tobacco exposure
- The panel testing contains, at minimum, testing of appropriate molecular aberrations (mutations, rearrangements, fusions, or amplifications) in **ALL** of the following genes: EGFR, ALK, ROS1, BRAF, ERBB2 (HER2), KRAS, MET, NTRK, and RET
- The individual is a candidate for targeted therapy that may be prescribed based on the panel test results
- The individual has not had prior NGS testing in the metastatic setting

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Metastatic non-small cell lung cancer (NSCLC) has historically been classified by histology and treated with cytotoxic chemotherapy. However, in the past decade there has been improved understanding of the disease biology and oncogenic driver mutations. The modern treatment of NSCLC is characterized by molecularly defined subsets actionable with targeted therapies and immune checkpoint inhibitors. It is estimated that 35%–50% of patients with advanced non-squamous NSCLC harbor a targetable alteration^{74, 75}, and selection of patients based on predictive biomarkers is associated with improved patient outcomes.⁷⁵ Molecular testing of sensitizing EGFR mutations, BRAF V600E, as well as ALK, ROS1, is now standard-of-care for patients with advanced NSCLC, as is testing for the EGFR T790M mutation upon resistance to first- or second generation EGFR tyrosine kinase inhibitor therapy.⁷⁶ Multiplexed genetic sequencing panels are preferred where available over multiple single gene tests to identify other treatment options beyond EGFR, ALK, BRAF, and ROS1. These larger panels include testing for specific molecular alterations in ERBB2 (HER2), KRAS, RET, MET, and NTRK genes.^{76, 77} New targetable alterations are continuing to emerge.

In contrast to metastatic nonsquamous NSCLC, the management of early-stage non-squamous NSCLC does not involve routine testing for oncogenic driver mutations. However, for patients with stage Ib to IIIA NSCLC who are being considered for FDA-approved adjuvant therapy with osimertinib (an oral targeted EFGR inhibitor) based on results of the ADAURA study⁷⁸, testing for EGFR exon 19 deletions or L858R point mutations is indicated. The ADAURA trial showed significant improvement in disease-free survival in these patients treated with adjuvant osimertinib; however, certain limitations of the design of this study raise questions about its role in adjuvant treatment. The overall survival data from the ADAURA trial and the results of another study, the ALCHEMIST trial, will influence practice patterns in the future.⁷⁹ Meanwhile, preclinical and retrospective clinical data that supports hypothesis testing of biomarker-driven treatment strategies in earlier stages of NSCLC with prospective clinical trials currently in progress.⁸⁰

Ovarian cancer (epithelial)

Somatic testing for pathogenic variants of BRCA1, BRCA2, and to determine HRD status in individuals with recurrent epithelial ovarian cancer is considered **medically necessary** when **ALL** of the following criteria are met:

- The individual has biopsy-proven epithelial ovarian cancer
- The individual is a candidate for treatment with a PARP inhibitor
- The individual has not had prior testing for these genes in the metastatic setting
- Testing involves an FDA-approved complementary diagnostic test used for ovarian cancer

Germline testing for pathogenic variants is considered **medically necessary** for all individuals with epithelial ovarian carcinoma. See *Hereditary Cancer Testing guideline* for further details.

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

The PARP inhibitor olaparib first received FDA approval in 2014 for women with germline mBRCA-associated (gBRCA) recurrent ovarian cancer. However, subsequent findings from clinical trials of PARP inhibitors have suggested that the importance of mBRCA as a predictive biomarker has diminished.⁸¹ For instance, among women without a germline mBRCA mutation, predictors of PARP activity include a somatic mutation in BRCA and evidence of homologous recombination deficiency. Indeed, FDA approval of the PARP inhibitor rucaparib in 2016 broadened the eligibility for PARP inhibitor treatment beyond gBRCA and included patients with somatic BRCA-associated ovarian cancers. In the subsequent Ariel3 trial, rucaparib significantly improved progression-free survival in patients with platinum-sensitive ovarian cancer who had achieved a

response to platinum-based chemotherapy.⁸² A novel aspect of this trial was the prospective validation of the tumor-based, NGS HRD test assay. Additionally, an exploratory analysis of progression-free survival in patients with BRCA wild-type and high loss of heterozygosity (LOH) or low LOH tumors revealed that patients with BRCA-mutant tumors did not solely drive rucaparib benefit in the HRD cohort or intention-to-treat population.⁸³ The FDA approval of rucaparib in April 2018 included approval of a complementary diagnostic test for tumor samples to determine both BRCA and HRD status.

Pancreatic Adenocarcinoma

Germline testing for pathogenic variants is considered **medically necessary** for all individuals with pancreatic adenocarcinoma. See *Hereditary Cancer Testing guideline* for further details.

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Up to 10% of individuals with pancreatic adenocarcinoma have a pathogenic germline alteration. Individuals with BRCA1 or BRCA2 mutations or microsatellite instability (MSI) resulting from a pathogenic germline mutation or somatic mutation in DNA mismatch repair (MMR) genes.⁸⁴ Affected individuals may benefit specifically from platinum-based therapies or PARP inhibitors.⁸⁵ For this reason, individuals with newly diagnosed pancreatic cancer at any stage should have multidisciplinary evaluation and management, germline testing, and integrated supportive care.⁸⁶

The enthusiasm for use of PARP-inhibition as targeted therapy for patients with BRCA-mutated prostate cancer came from the POLO trial published in 2019, a trial that randomized 154 patients with metastatic pancreatic adenocarcinoma and germline BRCA mutation to maintenance olaparib versus placebo after 16 weeks of first line platinum-based chemotherapy.⁸⁷ The POLO trial showed a statistically significant advantage in progression-free survival for olaparib (median 7.4 vs. 3.8 months, hazard ratio 0.53). However, mature data have since been published showing no statistical or clinically meaning improvement on overall survival.⁸⁸ Based on the absence of survival or quality of life improvement with olaparib and other concerns about the Polo trial design⁸⁹, enthusiasm for the use of olaparib as maintenance therapy has declined. A phase II trial is underway exploring the role of another PARP inhibitor, rucaparib, for maintenance therapy in patients with advanced pancreatic carcinoma and germline or somatic mutation of BRCA1, BRCA2, or PALB2.⁹⁰ These phase II data show that, like olaparib, rucaparib is active and tolerable. Ongoing randomized studies will further elucidate whether or not there are clinically meaningful benefits to using PARP inhibitors in the maintenance or adjuvant settings.

Prostate Cancer

Localized prostate cancer

Gene expression profiling and genomic biomarker tests as a technique for prostate cancer screening, detection, and management are considered **not medically necessary** for all indications.

Metastatic prostate cancer

Tissue-based NGS panel testing is considered **medically necessary** to identify pathogenic variants in individuals with metastatic prostate cancer when **ALL** of the following criteria are met:

- The individual has biopsy-proven adenocarcinoma of the prostate
- The individual is a candidate for **ONE** of the following therapies:
 - FDA-approved PARP inhibitor (olaparib, rucaparib, or another PARP inhibitor approved for use in this setting)
 - FDA-approved PD-1 inhibitor (pembrolizumab or another checkpoint inhibitor approved for use in this setting)
- The NGS panel includes BRCA2, BRCA1, and ATM, and may also include other genes encoding molecules involved in homologous recombination DNA repair such as PALB2, FANCA, RAD51D, CHEK2, and CDK12
- The individual has not had prior NGS testing in the metastatic setting

Germline testing for pathogenic variants is considered **medically necessary** for all individuals with metastatic prostate adenocarcinoma. See *Hereditary Cancer Testing guideline* for further details.

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Localized prostate cancer

Prostate cancer is usually suspected on the basis of a digital rectal exam (DRE) and/or an elevated prostate-specific antigen (PSA) test. Definitive diagnosis depends on histopathologic verification. Abnormal DRE is an indication for biopsy, but as an independent variable, PSA is a better predictor of cancer than either DRE or transrectal ultrasound.⁹¹ The histologic grading system for prostate cancer drives nearly all management decisions in localized prostate cancer, with Gleason score 6 being nearly universally indolent up to Gleason score 10 which is almost certainly lethal in the long run.⁹² The decision to proceed with a further staging workup is guided by which treatment options are available, taking into account the patient's preference and comorbidity. There are currently 10 or more pretreatment risk stratification tools for used in prostate cancer care, all of which use clinical and/or imaging factors without incorporating somatic genetic test information. The most commonly used are the D'Amico-derived systems (NCCN, NICE, GUROC, EAU, AUA) which involve categorization into 5 ordinal categories of risk: very low, low, intermediate, high, or very high. The Memorial Sloan Kettering nomogram, Cancer of the Prostate Risk score, and the Cambridge Prognostic Group are other systems and these perform slightly better in predicting prostate cancer death.⁹³ Prognostic approaches are sometimes explored using other, surrogate endpoints such as time to radiographic progression assessed by blinded independent central review, development of distant metastases, risk of adverse pathology during active surveillance, and others. Ultimately, management decisions for localized prostate cancer are typically made after appropriate options have been discussed with a multidisciplinary team (including urologists, radiation oncologists, medical oncologists, pathologists, and radiologists), and after the balance of benefits and side effects of each therapy modality has been considered in shared decision-making with the patient.

Numerous molecular biomarkers, particularly tissue-based genomic biomarker tests, have been developed to improve risk stratification and patient management. One of the unique challenges for use of these biomarkers is the complex spatial heterogeneity of prostate cancer.⁹⁴ While few of these genomic panels have undergone extensive validation, there are several commercially available tests (Oncotype DX prostate, Prolaris, Decipher, and ProMark) that have been shown in retrospective analyses to provide additional information beyond standard clinical models in prognostication or patient selection for therapy.^{95, 96} Given the absence of prospective clinical trial data, NCCN and ASCO guidelines do not recommend routine ordering of any molecular tests to guide decision-making in localized prostate cancer regarding the role of active surveillance or the use of post-prostatectomy adjuvant versus salvage radiation therapy. The ASCO guideline on molecular biomarkers in localized prostate cancer emphasizes that there is a paucity of prospective studies assessing the short and long-term outcomes of patients when these biomarkers are integrated into clinical decision-making.⁹⁵ These guidelines acknowledge that, based on lower level evidence and expert consensus, some specific molecular profiling biomarkers may be considered in specific situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. The most common settings where such testing is sometimes considered on that basis is with low or favorable intermediate-risk localized prostate cancer in men with life expectancy over 10 years. One limitation of the data regarding use of these tumor-tissue based genomic biomarker tests for active surveillance is that the tests were all developed in cohorts of patients who had already undergone primary treatment and were higher clinical risk than those typically considered for active surveillance.⁹⁷ Overall, it remains uncertain what prognostic endpoints should be prioritized and what magnitude of association with those prognostic endpoints are important. Nevertheless, in recent years, there has been more than 10-fold increase in tissue-based genomic biomarker testing related to prostate cancer with striking regional variability.⁹⁸ Practice patterns also vary widely within regions.⁹⁹ Issues surrounding clinician education and awareness of these assays (also referred to as "relationships with industry") may have contributed to this rising pattern of use.⁹⁷ The relative accuracy of these biomarker tests compared to other standard tests is unknown.⁹⁹ Also, while prospective trials are ongoing, the impact on key clinical outcomes (survival, quality of life, or need for treatment) attributable to use any of these tissue-based genomic biomarker tests (in any specific setting) is also uncertain.⁹⁷

Metastatic prostate cancer

Patients with metastatic prostate cancer have multiple treatment options with varied mechanisms of action beyond androgen deprivation therapy alone. Such options include androgen-receptor-targeted agents, taxane-based chemotherapies, bone-targeted radiopharmaceutical radium-223, and biomarker-driven therapy with the immune-checkpoint inhibitor pembrolizumab (for those with mismatch-repair deficiency (dMMR) or microsatellite instability (MSI)) and the PARP inhibitors olaparib and rucaparib (for those with homologous-recombination gene deficiency). Practice patterns vary in terms of the sequencing of therapies for both castrate-sensitive and castrate-resistant patients, and also variation in responses between patients with any given therapy.

The prevalence of recurrent genomic alternations varies across various prostate cancer clinical scenarios and also by published cohort. Common aberrations are typically not actionable and involve the androgen receptor (observed in >50% of cases); TP53 (in >40% of cases); genes encoding components of the PI3K pathway, such as PTEN (in 45% of cases); loss of RB1, which encodes the tumor suppressor Rb (in ~20% of cases); and others.¹⁰⁰ ESCAT level I molecular aberrations are those that the match of an alteration and a drug has been validated in clinical trials and should drive treatment decision in daily practice.³⁸ In this category, BRCA2, BRCA1, ATM and other genes encoding molecules involved in homologous recombination DNA repair (such as PALB2, FANCA, RAD51D, CHEK2, and CDK12) are found in 20%-25% of cases and may prompt consideration of PARP inhibitors. Moreover, roughly 3%–5% of prostate cancers harbor evidence of DNA mismatch-repair deficiency (dMMR), hyper-mutation or increased microsatellite instability which may prompt consideration of PD-1 inhibitors.¹⁰⁰ The ESMO Precision Medicine working group recommends multi-gene NGS panel testing in metastatic prostate cancer to assess for ESCAT level 1 alterations.³⁸ Commercially available prostate-cancer specific NGS panels include 11-14 genes.¹⁰¹ A metastatic biopsy for histologic and molecular evaluation is the standard of care and preferred over ctDNA testing, which can produce false positive biomarker signals due to potential interference from clonal hematopoiesis of indeterminate potential (CHIP). It is noteworthy that MSI-H status and HRD are generally mutually exclusive phenomena across cancer types, but may rarely co-occur, especially in prostate cancer. Most BRCA mutations coexisting with microsatellite instability are likely bystander events that may not result in sensitivity to poly (ADP-ribose) polymerase inhibitors.¹⁰²

The pivotal clinical trials of PARP inhibitors in metastatic castrate resistant prostate cancer are the phase III Profound trial (for olaparib) and the phase II Triton2 trial (for rucaparib). In the Profound trial^{103, 104}, there was a randomization to olaparib versus enzalutamide or abiraterone for patients who had either BRCA1, BRCA2, or ATM mutations (cohort A) or alterations in any of 12 other HRD genes (cohort B). The statistically significant benefit in progression free survival (7.4 months vs. 3.6 months, HR 0.34) and overall survival (19.1 vs. 14.7 months, HR 0.69) was limited to the cohort A patients. The phase II TRITON2 study of rucaparib included patients with mCRPC and deleterious BRCA or non-BRCA DNA damage-repair gene alterations treated after 1-2 lines of next-generation androgen-receptor directed therapy and 1 prior taxane-based regimen. In the BRCA mutated patients, the overall response rate was 43.5%¹⁰⁵, and for those with non-BRCA DNA damage-repair alterations the responses were much lower for PALB2, FANCA, BRIP1 and RAD51B and non-existent for ATM, CDK12, and CHEK2.¹⁰⁶

Thyroid Cancer

Testing of indeterminate thyroid nodules (ITN)

Use of next-generation gene expression classifier testing from fine needle aspirate sampling of a thyroid nodule is considered **medically necessary** when **ALL** of the following criteria are met:

- There has been no prior testing of the same thyroid nodule
- Initial cytopathology is reported as **ANY** of the following (Bethesda III or IV) categories:
 - Atypia of undetermined significance (AUS)
 - Follicular lesion of undetermined significance (FLUS)
 - Suspicious for follicular neoplasm (SFN)
 - Follicular neoplasm (FN)
- **ONE** of the following gene expression classifiers will be used:
 - Afirma Genomic Sequencing Classifier
 - ThyGeNEXT/ThyraMIR multiplatform test
 - ThyroSeq Genomic Classifier

Somatic genetic testing of thyroid malignancy

BRAF V600E testing is considered **medically necessary** for individuals with thyroid malignancy when the following criteria* are met:

- The individual has biopsy proven anaplastic thyroid cancer
- The individual is considered a potential candidate for FDA-approved oral targeted therapy based on the results of this testing

ALK, NTRK, and RET gene fusion testing is considered **medically necessary** for individuals with locally advanced, recurrent or metastatic thyroid carcinoma that is not amenable to radioactive iodine therapy when the following criteria are met:

- The individual has biopsy proven locally advanced, recurrent, or metastatic thyroid carcinoma
- The individual is considered a potential candidate for FDA-approved oral targeted therapy based on the results of this testing

**See additional guidelines concerning [tissue agnostic somatic testing](#) or [hereditary cancer risk testing](#) depending on the clinical scenario.*

Rationale

Molecular testing of indeterminate thyroid nodules

Roughly 4%-7% of the population have palpable thyroid nodules and as much as 30% or more having ultrasound detectable nodules. Indeterminate thyroid nodules (ITNs) are a challenging problem because these nodules are frequently discovered but usually benign and not requiring treatment.¹⁰⁷ The goal of care is to reduce overtreatment of such nodules and yet to detect and appropriately treat the proper subset of nodules that may cause morbidity or mortality due to thyroid malignancy. Fortunately, the majority of patients diagnosed with thyroid cancer are low risk for recurrence (<5%) and their risk of cancer-related death is even lower.¹⁰⁸ Indeed, recent studies have explored the role of de-escalation of treatment¹⁰⁸ including active surveillance.¹⁰⁹

When a thyroid nodule is found, the standard of care is to perform fine needle aspiration of the nodule for cytopathology. The American Thyroid Association (ATA) updated its guidelines for management of ITNs in 2015, recommending using the Bethesda System for Reporting Thyroid Cytopathology. This system includes six classifications: (I) nondiagnostic/unsatisfactory; (II) benign; (III) atypia of undetermined significance/follicular lesion of undetermined significance (AUS/FLUS); (IV) follicular neoplasm/suspicious for follicular neoplasm (FN); (V) suspicious for malignancy (SUSP); and (VI) malignant. Notably, the Bethesda System replaces the traditional single “indeterminate” category with three classifications, AUS/FLUS, FN, and SUSP.

Molecular testing is an increasingly utilized adjunct when evaluating indeterminate thyroid nodules in an effort to avoid unnecessary surgical or diagnostic risk to a patient.¹⁰⁷ The potential use of such tests is to improve risk stratification in the intermediate risk subgroups such as Bethesda III and IV groups. The second edition Bethesda System for Reporting Thyroid Cytology estimated 6%-18% malignancy rate of category III and 10%-40% for category IV nodules.¹¹⁰ A single institution retrospective cohort study noted in a 5-year review that malignancy rates at their institution of Bethesda III and IV surgical cases were 20%-25% with a fairly large range between years.¹¹¹ In recent years, several commercial multi-gene classifiers have emerged that are intended to help classify these indeterminate nodules.¹¹² One example is ThyroSeq v3, a third-generation DNA, and RNA sequencing test which identifies mutations, gene fusion, gene expression alterations, and copy number variations. In contrast, the Afirma GCS, another third-generation classifier, is a machine learning derived classification system detecting RNA transcriptome expression, mitochondrial RNA, and genomic copy number alterations including loss of heterozygosity. Finally, a multiplatform test (MPT) approach that combines a mutation panel test (ThyGenX) and a microRNA risk classifier test (ThyraMIR) has also been shown to provide both high NPV and high PPV for malignancy when tested in indeterminate nodules.^{113, 114} Of note, an altogether different classifier, the Afirma Xpression Atlas, detects gene variants and fusions in thyroid nodule FNA samples from a curated panel of 511 genes using whole-transcriptome RNA-sequencing. Its intended use is distinct from those above, focusing on cytologically indeterminate nodules that are Afirma GSC suspicious, Bethesda V/VI nodules, or known thyroid metastases.¹¹⁵

The ATA 2015 guidelines regarding evaluation and management of ITNs addressed the role of using molecular classifiers and biomarkers.¹¹⁶ The guidelines indicated that an ideal “rule-in” test would have a positive predictive value (PPV) for histopathologically proven malignancy similar to a malignant cytologic diagnosis (98.6%), and an ideal “rule-out” test would have a negative-predictive value similar to a benign cytologic diagnosis (96.3%). These guidelines commented on the limitations in the current literature, including lack of blinding of pathologists and clinicians, lack of controlled studies and direct comparisons of different evaluation methods, concerns about the reproducibility of findings across populations¹¹⁷, lack of clarity about how test results change surgical management, and lack of long-term outcome data. Overall, these ATA guidelines were praised by community oncologists and European specialists for more strongly emphasizing nodule and tumor morphology and topography and for showing less enthusiasm for molecular biomarkers for risk stratification (compared to their prior guideline), emphasizing the relative importance of clear, comprehensive ultrasonographic and pathology reports.¹¹⁸ Molecular testing is thus viewed as an adjunct that may facilitate decision-making in selected patients. The ATA guideline rationales are that no single molecular test or combination of tests yet has proved useful in terms of clinical utility across all indeterminate cases and that molecular testing continues to evolve.¹¹⁹ Subsequent studies, including a large, blinded, multicenter study of Thyroseq v3 showed that the testing did not meet the ATA threshold for a “rule-in” test due to the low PPV and its potential value as a rule-out test varied by cancer prevalence but was sufficiently high (NPV 97%) when that

prevalence was 28%.¹²⁰ A systematic review and meta-analysis of published gene expression classifier studies in this realm observed that the published validation cohorts were not representative of the populations in whom these tests have been used, particularly because of differences in the underlying prevalence rates of cancer which affect the negative predictive value of the test.¹²¹

Overall, several molecular classifiers have demonstrated analytical and clinical validity in the evaluation of ITNs. However, for any biomarker test to have clinical utility, it must also be shown to improve outcomes compared with if the biomarker test results are not known.¹²² The prevalence of thyroid cancer in these nodules varies considerably between studies and between sites in the multicenter studies.^{121, 123} Thus, to apply these molecular test results in patient management, clinicians must know the prevalence of cancer in their own patient population where the test would be applied. Furthermore, clinicians must be careful in deciding when to use the test, because for patients with an indication for surgery based on cytology alone (or contraindication for surgery based on various reasons), there is little value in pursuing this type of testing.¹²³

There is special interest in ITNs with Hürthle cells. Hürthle cells are thyroid follicular-derived epithelial cells with oncocytic cytology, and accurately classifying nodules with these cells is difficult with fine-needle aspiration. There has been interest in the use of third generation molecular classifiers for this subset of ITNs. Although patients with advanced Hürthle cell carcinoma have a poor prognosis, patients presenting with minimally invasive, early disease have an excellent prognosis.¹²⁴ The majority of these lesions have low-risk or no molecular alterations and are benign on follow-up. Unfortunately, no single molecular alteration defines cytologically indeterminate Hürthle cell lesions; and molecular testing is not definitive in determining which cases may be managed conservatively.¹²⁵ There are ongoing efforts to improve classification of these nodules¹²⁶, but thus far the accuracy of these molecular classifier tests falls short of providing clinical utility to guide surgical management.

Unknown Primary Site Cancer

Somatic genetic testing for individuals with cancer of unknown primary site is considered **not medically necessary**.

Note: Tumor agnostic genetic testing indications may also apply, depending on the clinical scenario. See the [Tumor Agnostic Testing](#) guideline for details.

Rationale

Cancer of unknown primary site (CUP) refers to a diverse set of malignancies where the tissue of origin remains unclear the standard set of detailed investigations (clinic, laboratory, pathology, and imaging) have been completed. These may also be called occult primary cancers. With the advent of modern imaging and immunohistochemistry diagnostic techniques, the incidence of CUP has decreased from around 3%–5% in the 1990s to 1%–2% in the current era.¹²⁷ CUP is categorized into four histological types that include adenocarcinoma of good-to-moderate differentiation (50%), followed by poorly undifferentiated adenocarcinomas (30%), squamous cell carcinoma (15%) and undifferentiated neoplasms (5%).¹²⁸ In most patients, the disease is disseminated and incurable. The majority of patients present with visceral disease, mainly to liver and lungs. CUP was once treated as a distinct cancer and investigated with phase 2 treatment trials response rates of 25% to 35% and survival ranging from 6 to 16 months. Survival has been longer for patients with nodal, pleural, or serous peritoneal disease (14 to 16 months) than for patients with visceral metastatic disease (6 to 9 months).¹²⁹ Less common, presentations such as bone predominant CUP (with median survival around 15 months) or lymph node-only CUP (median survival over 30 months) drive unique approaches in terms of their risk stratification and treatment.¹³⁰ In a broad phase II trial using pembrolizumab to treat rare cancers, 22 patients with CUP were enrolled with no complete responses, 3 partial responses, and 4 patients with stable disease.¹³¹

Various tissue of origin (TOO) classifiers have been developed TOO classifiers have been developed on data from a wide range of molecular methods including targeted DNA sequencing, whole exome and whole genome DNA sequencing, RNA profiling, and methylation profiling.¹³² Several whole genome sequencing (WGS) based have been developed and are able to discriminate between 18 and 35 cancer types but with different performance across sub-cohorts of common cancer types (such as breast and colorectal cancer) compared to various other types of uncommon cancers.¹³²⁻¹³⁴ There is also some variation based on TOO classifiers that are testing untreated versus treated metastatic cancers.¹³² Unfortunately, advances in diagnostics have not yielded clinical utility as yet through translation into a survival benefit, as no differences in outcome were reported between empirical and molecularly guided treatments.¹²⁸ Meanwhile, user-friendly nomograms using readily available clinicopathological factors (not based on molecular testing) can provide robust personalized prognostication and aid in decision-making and selection or stratification of patients for clinical trial enrollment.¹³⁵

Somatic Testing of Hematologic Malignancies

Clinical Indications

General Requirements

Repeated testing of the same individual for the same indication should be limited to evaluation following an intervention, or when there is a change in clinical status such that additional testing is required to determine next steps in management. At times, it may be necessary to repeat a test using different techniques or protocols to clarify a finding or result of the original study.

Repeated testing for the same indication using the same or similar technology may be subject to additional review or require peer-to-peer conversation in the following scenarios:

- Repeated diagnostic testing of the same tumor site with no clinical change, treatment, or intervention since the previous study
- Repeated diagnostic testing of the same individual and the same tumor by different providers for the same member over a short period of time

Umbrella Criteria

Somatic Genomic Testing (Tumor Biomarker Testing)

Somatic genomic testing is considered **medically necessary** in individuals with cancer when **ANY** of the following criteria are met:

- The genomic testing has established analytical and clinical validity and is performed in an appropriately certified laboratory
- The genetic test has established clinical utility such that a positive or negative result will meaningfully impact the clinical management (predictive, diagnostic, prognostic, or therapeutic) of the individual and will likely result in improvement in net health outcomes (i.e., the health benefits of the interventions outweigh any medical or psychological harmful effects of the testing intervention)
 - When there are genomic biomarker-linked therapies approved by the U.S. Food and Drug Administration (FDA) for the individual's specific cancer scenario and such therapies are being considered in the near term
 - When considering a treatment for which there are specific genomic biomarker-based contraindications or exclusions related therapeutic decisions being considered in the near term
- Clinical decision making incorporates the known or predicted impact of a specific genomic alteration on protein expression or function and published clinical data on the efficacy of targeting that genomic alteration with a particular agent

Rationale

Nearly every malignancy will have somatic mutations that have been described, although most known mutations do not have clinical management implications. While various common conditions are covered by specific guideline criteria for somatic testing of tumors, it is not feasible to establish criteria for every clinical scenario in oncology and hematology. The general criteria for somatic testing (above) apply to malignancy when more specific criteria are not available.

Cancer-specific Criteria

Acute Lymphocytic Leukemia

Tissue-based somatic genetic testing is considered **medically necessary** for children and adults with acute lymphoblastic leukemia to establish the diagnosis or to identify actionable therapeutic targets when **ANY** of the following criteria are met:

- A multi-gene panel contains, at a minimum, the following genes: ABL1, ABL2, CRLF2, CSF1R, FLT3, IL7R, JAK1, JAK2, JAK3, PDGFRB, SH2B3, TP53, and IKZF1

Chromosomal analyses of bone marrow specimens, which may also include FISH testing, to detect and characterize clonal chromosomal abnormalities that have important diagnostic, prognostic, and therapeutic implications are considered **medically necessary** for children and adults with acute lymphoblastic leukemia.

The use of NGS testing of tumor DNA is considered **medically necessary** to detect or quantify minimal residual disease in children or adults with acute lymphocytic leukemia.

BCR-ABL kinase domain point mutation analysis is considered **medically necessary** in the evaluation of individuals with BCR-ABL (Philadelphia chromosome) positive acute lymphoblastic leukemia to evaluate treated individuals who manifest suboptimal response to initial tyrosine kinase inhibitor therapy or loss of response to tyrosine kinase inhibitor therapy.

PCR testing for BCR-ABL1 quantification is considered **medically necessary** in the monitoring of Philadelphia chromosome-positive ALL.

Rationale

Acute lymphoblastic leukemia (ALL) has a bimodal age distribution, the first peak occurring at approximately 5 years of age and the second at approximately 50 years. ALL may be of B-cell precursor or T-cell lineage. In the United States, ALL is the most common cancer among children and the most frequent cause of death from cancer before 20 years of age.¹³⁶ The cure rate of pediatric ALL is at least 80%, in adults closer to 30% to 40%.¹³⁷ Several genetic factors (such as Down's syndrome) are associated with an increased risk of ALL, but most patients have no recognized inherited factors.

On average, childhood ALL genomes contain 10 to 20 non-silent coding mutations at the time of diagnosis and about twice as many at the time of relapse. Many mutations affect the transcriptional regulation of lymphoid development and differentiation; cell-cycle regulation; the TP53–retinoblastoma protein tumor-suppressor pathway; growth factor receptor, Ras, phosphatidylinositol 3-kinase, and JAK-STAT signaling; nucleoside metabolism; and epigenetic modification.¹³⁶ Adult patients have been found to have more gene mutations, especially IKZF1, MLL2, and JAK3, but fewer alterations of PTPN11 compared with pediatric patients.¹³⁸ The precise contribution of these genetic or epigenetic abnormalities to leukemogenesis, the development of drug resistance and leukemic clone evolution remains to be defined.¹³⁸

Quantification of measurable residual disease (MRD) detected by quantitative polymerase chain reaction (PCR), flow cytometry, or next generation sequencing (NGS). The risk of ALL treatment failure and death is 3 to 5 times as high among children with levels of minimal residual disease that are 0.01% or higher at the end of induction therapy and at later time points than among those with levels that are lower than 0.01%.¹³⁶ Overall, MRD testing has prognostic implications in various subsets of both pediatric and adult ALL across therapies, methods of and times of MRD assessment, cutoff levels, and disease subtypes based on a meta-analysis of 39 publications comprising over 13,000 patients.¹³⁹ There are caveats with this testing. Although MRD is a direct measure of disease burden and treatment response in ALL, there may be sanctuary sites in the body that contribute to relapse but are not measurable by conventional methods. Also, technical difficulties can give misleading results, and standardized methods for MRD determination are not widely available outside of specialized centers.¹³⁹ A randomized clinical trial of intensification of therapy for patients with higher levels of minimal residual disease has been conducted showing non-significant improvement in survival, but demonstrating the promise of this method of patient selection for clinical trials.¹⁴⁰ MRD monitoring has become integral to clinical trials developed by the St. Jude's Consortium, the Children's Oncology Group, and the Dana Farber Cancer Institute Consortium and also recognized as an essential component of patient evaluation and over the course of sequential therapy in consensus-based guidelines.¹⁴¹

Acute Myelogenous Leukemia

Tissue-based somatic genetic testing is considered **medically necessary** for individuals with acute myelogenous leukemia (AML) to establish the diagnosis and to identify actionable therapeutic targets when **ANY** of the following criteria are met:

- A multi-gene panel contains, at a minimum, the following genes: FLT3, IDH1, IDH2, NPM1, CEBPA, DDX41, TP53; ASXL1, BCOR, EZH2, RUNX1, SF3B1, SRSF2, STAG2, U2AF1, ZRSR2

Chromosomal analyses of bone marrow specimens, which may also include FISH testing, to detect and characterize clonal chromosomal abnormalities that have important diagnostic, prognostic, and therapeutic implications are considered **medically necessary** for individuals with AML.

Rationale

Acute myelogenous leukemia (AML) is a disease characterized by infiltration of the bone marrow, blood, and other tissues by proliferative, clonal, abnormally differentiated hematopoietic cells. Most cases of AML are characterized by clonal heterogeneity at the time of diagnosis. Leukemic relapse invariably arises from a pre-existing—or at least a closely genetically-related—clone.¹⁴² Various patterns of dynamic clonal evolution that occur at relapse probably contribute to resistance to therapy.¹⁴³ Adults under age 60 years are cured with intensive leukemia treatment around 40% of the time.¹⁴³ However, this is generally a disease of older adults with a median age at diagnosis of 68 years and escalating incidence with increasing age, with older adults having a much worse prognosis and cure rate closer to 10%.¹⁴⁴

The World Health Organization (WHO) 5th edition was updated in 2022 and the classification of AML emphasizes recent breakthroughs in how this disease is understood and managed. Foremost is the separation of AML with defining genetic abnormalities from AML defined by differentiation.¹⁴⁵ The defining genetic abnormalities include the following: fusions of PML::RARA, RUNX1::RUNX1T1CBFB::MYH11, DEK::NUP214, RBM15::MRTFA, and BCR::ABL1, rearrangements of KMT2A, MECOM, NUP98, NPM1, or CEBPA, and mutations of NPM1 or CEBPA. Genomic evaluation of patients with AML has been demonstrated to have a higher diagnostic yield compared to conventional cytogenetic analysis.¹⁴⁶

The most widely used consensus risk stratification guidelines in AML are those from the European LeukemiaNet (ELN), an international working group of several dozen AML experts. The 2022 ELN recommendations are for screening for the following specific gene mutations required for establishing the diagnosis and to identify actionable therapeutic targets: FLT3, IDH1, IDH2, NPM1, CEBPA, DDX41, TP53; ASXL1, BCOR, EZH2, RUNX1, SF3B1, SRSF2, STAG2, U2AF1, ZRSR2.¹⁴⁷ The ELN acknowledges that additional genes are investigational in this rapidly evolving field and are not currently required for establishing the diagnosis or for the identification of actionable therapeutic targets. Leukemia experts have advocated for the principle that risk groups are dynamic as newer therapies emerge for specific disease subsets and updated approaches may be recognized and widely adopted by nationally and internationally.¹⁴⁸

Measurable residual disease (MRD) can be detected by polymerase chain reaction (PCR), flow cytometry, or next generation sequencing (NGS). MRD testing has prognostic implications in various subsets of AML and at various times throughout treatment.^{149, 150} This testing may define molecular persistence after intensive chemotherapy, molecular progression or relapse after intensive chemotherapy, or molecular relapse, progression, or persistence post stem cell transplantation.¹⁵¹ Testing is often pursued to assess eligibility in MRD-directed clinical trials. The proven therapeutic implications of MRD remain mostly unresolved.^{142, 147}

Chronic Myeloid Leukemia and Myeloproliferative Disorders

Bone marrow tissue-based somatic genetic testing is considered **medically necessary** for establishing the diagnosis of a myeloproliferative disorder when **ANY** of the following criteria are met:

- PCR or FISH testing of BCR-ABL1 is performed to evaluate for chronic myeloid leukemia
- PCR, FISH, or NGS testing of JAK2, CALR, and MPL is performed to evaluate for BCR-ABL1 (Philadelphia chromosome) negative myeloproliferative disorders

BCR-ABL kinase domain point mutation analysis is considered **medically necessary** in the monitoring of chronic myeloid leukemia (CML) in **ANY** of the following circumstances:

- Evaluation of individuals with chronic myelogenous leukemia to evaluate treated individuals who manifest suboptimal response to tyrosine kinase inhibitor therapy indicated by:
 - Lack of a partial hematologic or cytogenetic response at 3 months or greater after treatment onset

- Less than a complete hematologic and cytogenetic response at 12 months
- Disease progression to accelerated or blast phase

Chromosomal analyses of bone marrow specimens to detect and characterize clonal chromosomal abnormalities that have important diagnostic, prognostic, and therapeutic implications are considered **medically necessary** for individuals with chronic myeloid leukemia.

PCR testing for BCR-ABL1 quantification is considered **medically necessary** for response assessment every 3 months during active treatment and every 6 weeks in the first year after treatment discontinuation.

Rationale

Elevated peripheral blood cell counts, such as leukocytosis, thrombocytosis, and polycythemia, are often the presenting symptom in patients with myeloproliferative neoplasms (MPN). These diseases include chronic myeloid leukemia (CML), polycythemia vera (PV), essential thrombocytosis (ET), primary myelofibrosis (PM), and several other less common subtypes.¹⁴⁵ PMNs are characterized by the abnormal proliferation of mature bone marrow cell lineages and a clinical course that is usually prolonged, but ultimately leads to marrow fibrosis and acute leukemia. These are hematopoietic stem-cell disorders that share mutations that constitutively activate the physiologic signal-transduction pathways responsible for hematopoiesis.¹⁵² Likewise, CML is a disease of hematopoietic stem cells arising from a translocation t(9;22)(q34;q11) with the shortened chromosome 22. The translocation leads to a juxtaposition of the ABL1 gene from chromosome 9 and the BCR gene from chromosome 22, resulting in a BCR-ABL1 fusion gene that codes for BCR-ABL1 transcripts and fusion proteins with high tyrosine kinase activity.¹⁵³ Like MPN, patients with CML typically present with elevated peripheral blood counts, particularly excessive granulocytes and a left shift of granulopoiesis, and common signs and symptoms include fatigue, weight loss and splenomegaly.

Patients with signs and symptoms consistent with CML or MPN are typically evaluated with review of the peripheral blood smear review and BCR-ABL1 analysis undertaken for consideration of CML. In the absence of BCR-ABL1 translocation and lack of dysplasia, molecular characterization of JAK2, CALR, and MPL are pursued as well as bone marrow morphologic review for an accurate diagnosis.¹⁵⁴ A positive finding for a mutation established the presence of a hematopoietic stem cell disorder and rules out myelodysplasia, but none of these mutations are disease-defining.¹⁵² Distinction between the MPN types is based on integrating peripheral blood findings with molecular data and bone marrow morphologic evaluation findings, as none of these parameters alone provide sufficient diagnostic specificity.¹⁴⁵ Several other somatic mutations have been reported in persons with JAK2, MPL or CALR mutations including epigenetic modifiers (ASXL1, TET2, EZH2, IDH1, IDH2, DNMT3A), RNA splicing factors (SRSF2, U2AF1, SF3B1) and transcriptional regulators (TP53, IKZF1, NF-E2, CUX1). These mutations do not cause myeloproliferation.¹⁵⁵ The cytogenetic landscape of MPNs is limited and does not differ substantially according to the type of neoplasm. Moreover, driver mutation status is not associated with the time to leukemic transformation or survival after transformation.¹⁵²

For CML specifically, the diagnosis must be confirmed by cytogenetics showing t(9;22)(q34;q11) and by multiplex RT-PCR showing BCR-ABL1 transcripts. At baseline, neither quantitative RT-PCR measuring BCR-ABL1 transcripts level nor BCR-ABL1 mutation analysis are advised at baseline since these data are not necessary for decision-making.¹⁵³ Tyrosine kinase inhibitor therapy (such as imatinib and other agents) and careful disease monitoring has reduced the incidence of progression to advanced phase and the 10-year overall survival rate for CML is now 80%-90%.¹⁵⁶ A principal goal of CML therapy with TKIs is the achievement of a stable molecular remission which accords successful discontinuation of therapy, a concept now known as 'treatment-free remission'.¹⁵⁵ Monitoring of treatment is now almost exclusively done molecularly by quantitative PCR measurement of BCR-ABL1 transcripts according to the International Scale with optimal response, warning category, and treatment failure defined.¹⁵⁷ BCR-ABL1 $\leq 1\%$ has been determined to be equivalent to complete cytogenetic remission; greater than 10% BCR-ABL1 at 3 months indicates treatment failure when confirmed. Response milestones are the same for first and second-line therapy and include measurements at baseline and every 3 months thereafter with more intense monitoring (every 4-6 weeks) for the first year after treatment discontinuation.¹⁵³ Achieving specific time-dependent molecular milestones, as defined by global therapeutic guidelines, has been established as critical in maximizing optimal outcomes while identifying patients at risk of therapy failure.¹⁵⁸

More than 100 different kinase domain mutations of BCR-ABL1 that impair TKI binding have been reported in patients who develop TKI resistance.¹⁵⁶ Resistance to imatinib occurs in 10%–15%, and to second generation TKIs in <10% of patients in first-line treatment. Mutations account for resistance in about one-third of resistant patients in chronic phase, and in about two-thirds of resistant patients in accelerated or blast phase.¹⁵⁹ A prospective multicenter study has demonstrated that NGS provides a more accurate picture of BCR-ABL1 mutation status compared to standard Sanger sequencing and demonstrated the clinical relevance of low level mutations¹⁶⁰, thus NGS is the recommended technology for detecting BCR-ABL1 resistance mutations in patients not adequately responding to TKI therapy.¹⁵⁹

Multiple Myeloma

Gene expression profile tests

Gene expression profile tests for diagnostic evaluation, risk stratification, or management of multiple myeloma are considered **not medically necessary**.

Chromosomal analyses of bone marrow specimens

Chromosomal analyses of bone marrow specimens to detect and characterize clonal chromosomal abnormalities that have important diagnostic, prognostic, and therapeutic implications are considered **medically necessary** for individuals with multiple myeloma.

The use of NGS testing of tumor DNA to detect or quantify minimal residual disease (MRD) in individuals with myeloma is considered **medically necessary** under **EITHER** of the following circumstances:

- MRD testing used prior to initiating new treatment intended to induce myeloma remission
- MRD testing used to assess depth of response after a cycle of treatment intended to induce myeloma remission

Rationale

Multiple myeloma is the second most common hematological malignancy, accounting for about 2% of cancer deaths in the United States. Almost all patients with multiple myeloma evolve from an asymptomatic pre-malignant stage termed monoclonal gammopathy of undetermined significance (MGUS), and the rate of progression is influenced by the underlying cytogenetic findings.¹⁶¹ The diagnosis requires the presence of clonal plasma cells in the bone marrow or in a biopsy-proven bone or extramedullary plasmacytoma. The majority of patients with myeloma present with symptoms related to organ involvement (hypercalcemia, renal insufficiency, anemia, and bone lesions). Less commonly, patients may be diagnosed based on abnormal blood or urine tests.¹⁶²

Multiple myeloma is increasingly recognized as more than one disease, characterized by marked cytogenetic, molecular, and proliferative heterogeneity. Risk stratification is important for prognostication, patient selection for clinical trials, and comparison of treatment approaches. Risk stratification is complex and involves cytogenetic classification combined with disease stage, host factors, and other prognostic factors.¹⁶¹ The revised ISS(R-ISS) was introduced in 2015 and has been widely validated.¹⁶³ More recently, additional cytogenetic abnormalities have been found and newer, validated models are available such as the Mayo Additive Staging System¹⁶⁴ and evolving tools such as the Mayo Clinic mSMART risk stratification method (www.msmaart.org).¹⁶¹

Given the high rates of complete response seen in patients with multiple myeloma with new treatment approaches, new response categories have emerged that focus on detection of residual tumor cells in the bone marrow using modern flow cytometry or next-generation sequencing (NGS) techniques. The clinical utility of peripheral blood-based evaluation of minimal residual disease remains unknown.¹⁶⁵ MRD negativity in the bone marrow is defined as the absence of tumor plasma cells within 1,000,000 bone marrow cells, a threshold that indicates the best results for the prediction of both progression-free and overall survival.¹⁶⁶ Detailed aspects of this approach to detecting minimal residual disease (MRD) were outlined in 2016 by the International Myeloma Working Group (IMWG)¹⁶⁵, and guidelines from the American Society and Cancer Care Ontario¹⁶² and from the European Society of Medical Oncology and the European Hematology Association¹⁶⁶ have since endorsed routine use of MRD testing for response assessment. The ASCO guideline makes a strong recommendation that the quality and depth of response should be assessed by IMWG criteria, with the recommendation based on high-quality evidence. MRD is recommended to test the depth of response after each cycle of therapy intended to induce remission. The use of MRD to assess disease status during maintenance therapy is still unknown. For example, it is unclear whether maintenance therapy in MRD-negative patients can be stopped or whether treatment needs to be changed in MRD-positive patients.¹⁶⁶ ASCO considers the evidence strength low and recommendation weak for MRD testing during maintenance therapy, and makes a moderate strength recommendation not to modify maintenance therapy based on depth of response information.¹⁶²

Myelodysplastic Syndrome

Somatic testing of bone marrow tissue is considered **medically necessary** for individuals with clinically diagnosed or suspected myelodysplastic syndrome when **ANY** of the following criteria are met:

- Testing is for the purpose of establishing the diagnosis or to identify actionable therapeutic targets

- A targeted multi-gene panel contains, at a minimum, the following genes: ASXL1, DNMT3A, EZH2, NRAS, RUNX1, SF3B1, SRSF2, STAG2, TET2, TP53, U2AF1, ZRSR2

Chromosomal analyses of bone marrow specimens to detect and characterize clonal chromosomal abnormalities that have important diagnostic, prognostic, and therapeutic implications are considered **medically necessary** for individuals with myelodysplastic syndrome.

Rationale

Myelodysplastic syndromes (MDS) are clonal hematopoietic neoplasms defined by cytopenias and morphologic dysplasia. MDS occurs mainly, but not exclusively in older adults, with the median age being 70 years. MDS evolves to AML through a process of clonal selection, with the pattern of transformation occurring in different patterns depending on the subtype and mutation drivers.¹⁶⁷ In subtypes of MDS at lower risk of transformation, treatment is focused mainly on improving anemia and other cytopenias. In higher risk disease, the focus is more on delaying disease progression and prolonging survival. Allogeneic stem cell transplantation is the only potentially curative therapy, but it is rarely applicable since most patients are older adults.¹⁶⁸ The World Health Organization (WHO) 5th edition was updated in 2022 and the classification of MDS now features genetically defined disease types rather than risk-based grouping based on features such as blast percentage, ring sideroblasts, and number of lineages with dysplasia.¹⁴⁵ Somatic mutations with an overall incidence of 5% or greater in MDS include ASXL1, DNMT3A, EZH2, NRAS, RUNX1, SF3B1, SRSF2, STAG2, TET2, TP53, U2AF1, and ZRSR2.¹⁶⁹

To meet the challenge of the heterogeneity in clinical course and outcomes for MDS, the International Prognostic Scoring System–Revised (IPSS-R), was developed by the International Working Group for Prognosis in MDS and set the course for risk stratification, clinical trial design, and treatment recommendations.¹⁷⁰ More recently, a clinical-molecular prognostic model (IPSS-Molecular [IPSS-M]) has been validated.¹⁷¹ In multivariable analysis, the top genetic mutations found to be predictors of adverse outcomes were identified in TP53, FLT3, and KMT2A (MLL), while SF3B1 mutations were associated with favorable outcomes. Moreover, mutations in ASXL1, BCOR, EZH2, NRAS, RUNX1, STAG2, and U2AF1 were significantly associated with adverse risk for several key outcomes as well.

References

1. Kurnit KC, Dumbrava EEI, Litzenger B, et al. Precision Oncology Decision Support: Current Approaches and Strategies for the Future. *Clin Cancer Res*. 2018;24(12):2719-31. Epub 2018/02/09. PMID: 29420224
2. Chakravarty D, Johnson A, Sklar J, et al. Somatic Genomic Testing in Patients With Metastatic or Advanced Cancer: ASCO Provisional Clinical Opinion. *J Clin Oncol*. 2022;40(11):1231-58. Epub 2022/02/18. PMID: 35175857
3. Cobain EF, Wu YM, Vats P, et al. Assessment of Clinical Benefit of Integrative Genomic Profiling in Advanced Solid Tumors. *JAMA Oncol*. 2021;7(4):525-33. Epub 2021/02/26. PMID: 33630025
4. Le Tourneau C, Delord JP, Gonçalves A, et al. Molecularly targeted therapy based on tumour molecular profiling versus conventional therapy for advanced cancer (SHIVA): a multicentre, open-label, proof-of-concept, randomised, controlled phase 2 trial. *Lancet Oncol*. 2015;16(13):1324-34. Epub 2015/09/08. PMID: 26342236
5. El-Deiry WS, Goldberg RM, Lenz HJ, et al. The current state of molecular testing in the treatment of patients with solid tumors, 2019. *CA Cancer J Clin*. 2019;69(4):305-43. Epub 2019/05/23. PMID: 31116423
6. André T, Cohen R, Salem ME. Immune Checkpoint Blockade Therapy in Patients With Colorectal Cancer Harboring Microsatellite Instability/Mismatch Repair Deficiency in 2022. *Am Soc Clin Oncol Educ Book*. 2022;42:1-9. Epub 2022/04/27. PMID: 35471834
7. McGrail DJ, Pilié PG, Rashid NU, et al. High tumor mutation burden fails to predict immune checkpoint blockade response across all cancer types. *Ann Oncol*. 2021;32(5):661-72. Epub 2021/03/20. PMID: 33736924
8. Jardim DL, Goodman A, de Melo Gagliato D, et al. The Challenges of Tumor Mutational Burden as an Immunotherapy Biomarker. *Cancer cell*. 2021;39(2):154-73. Epub 2020/10/31. PMID: 33125859
9. Zehir A, Benayed R, Shah RH, et al. Mutational landscape of metastatic cancer revealed from prospective clinical sequencing of 10,000 patients. *Nat Med*. 2017;23(6):703-13. Epub 2017/05/10. PMID: 28481359
10. Sung MT, Wang YH, Li CF. Open the Technical Black Box of Tumor Mutational Burden (TMB): Factors Affecting Harmonization and Standardization of Panel-Based TMB. *Int J Mol Sci*. 2022;23(9). Epub 2022/05/15. PMID: 35563486
11. Subbiah V, Solit DB, Chan TA, et al. The FDA approval of pembrolizumab for adult and pediatric patients with tumor mutational burden (TMB) ≥ 10 : a decision centered on empowering patients and their physicians. *Ann Oncol*. 2020;31(9):1115-8. Epub 2020/08/11. PMID: 32771306
12. Prasad V, Addeo A. The FDA approval of pembrolizumab for patients with TMB > 10 mut/Mb: was it a wise decision? No. *Ann Oncol*. 2020;31(9):1112-4. Epub 2020/08/11. PMID: 32771305

13. Solomon JP, Benayed R, Hechtman JF, et al. Identifying patients with NTRK fusion cancer. *Ann Oncol.* 2019;30(Suppl_8):viii16-viii22. Epub 2019/11/19. PMID: 31738428
14. Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol.* 2020;21(2):271-82. Epub 2019/12/16. PMID: 31838007
15. Drilon A, Laetsch TW, Kummar S, et al. Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children. *N Engl J Med.* 2018;378(8):731-9. Epub 2018/02/22. PMID: 29466156
16. Solomon JP, Linkov I, Rosado A, et al. NTRK fusion detection across multiple assays and 33,997 cases: diagnostic implications and pitfalls. *Mod Pathol.* 2020;33(1):38-46. Epub 2019/08/04. PMID: 31375766
17. Mateo J, Chakravarty D, Dienstmann R, et al. A framework to rank genomic alterations as targets for cancer precision medicine: the ESMO Scale for Clinical Actionability of molecular Targets (ESCAT). *Ann Oncol.* 2018;29(9):1895-902. Epub 2018/08/24. PMID: 30137196
18. Hilbers FS, Aftimos P. Expanding the landscape of actionable genomic alterations in metastatic breast cancer: comprehensive genomic profiling for all? *Ann Oncol.* 2020;31(8):967-9. Epub 2020/05/19. PMID: 32422170
19. Subbiah V, Baik C, Kirkwood JM. Clinical Development of BRAF plus MEK Inhibitor Combinations. *Trends cancer.* 2020;6(9):797-810. Epub 2020/06/17. PMID: 32540454
20. Subbiah V, Kreitman RJ, Wainberg ZA, et al. Dabrafenib plus trametinib in patients with BRAF V600E-mutant anaplastic thyroid cancer: updated analysis from the phase II ROAR basket study. *Ann Oncol.* 2022;33(4):406-15. Epub 2022/01/14. PMID: 35026411
21. Wen PY, Stein A, van den Bent M, et al. Dabrafenib plus trametinib in patients with BRAF(V600E)-mutant low-grade and high-grade glioma (ROAR): a multicentre, open-label, single-arm, phase 2, basket trial. *Lancet Oncol.* 2022;23(1):53-64. Epub 2021/11/29. PMID: 34838156
22. Subbiah V, Lassen U, Élez E, et al. Dabrafenib plus trametinib in patients with BRAF(V600E)-mutated biliary tract cancer (ROAR): a phase 2, open-label, single-arm, multicentre basket trial. *Lancet Oncol.* 2020;21(9):1234-43. Epub 2020/08/21. PMID: 32818466
23. Matulay JT, Kamat AM. Advances in risk stratification of bladder cancer to guide personalized medicine. *F1000Res.* 2018;7. Epub 2018/08/16. PMID: 30109022
24. Robertson AG, Kim J, Al-Ahmadie H, et al. Comprehensive Molecular Characterization of Muscle-Invasive Bladder Cancer. *Cell.* 2017;171(3):540-56.e25. Epub 2017/10/11. PMID: 28988769
25. Warrick JI, Knowles MA, Yves A, et al. Report From the International Society of Urological Pathology (ISUP) Consultation Conference On Molecular Pathology Of Urogenital Cancers. II. Molecular Pathology of Bladder Cancer: Progress and Challenges. *Am J Surg Pathol.* 2020;44(7):e30-e46. Epub 2020/02/25. PMID: 32091435
26. Lenis AT, Lec PM, Chamie K, et al. Bladder Cancer: A Review. *Jama.* 2020;324(19):1980-91. Epub 2020/11/18. PMID: 33201207
27. Loriot Y, Necchi A, Park SH, et al. Erdafitinib in Locally Advanced or Metastatic Urothelial Carcinoma. *N Engl J Med.* 2019;381(4):338-48. Epub 2019/07/25. PMID: 31340094
28. Necchi A, Madison R, Pal SK, et al. Comprehensive Genomic Profiling of Upper-tract and Bladder Urothelial Carcinoma. *Eur Urol Focus.* 2021;7(6):1339-46. Epub 2020/08/31. PMID: 32861617
29. Bhutiani N, Egger ME, Ajkay N, et al. Multigene Signature Panels and Breast Cancer Therapy: Patterns of Use and Impact on Clinical Decision Making. *J Am Coll Surg.* 2018;226(4):406-12.e1. Epub 2018/01/26. PMID: 29366844
30. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2018;379(2):111-21. Epub 2018/06/05. PMID: 29860917
31. Sparano JA, Gray RJ, Makower DF, et al. Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2015;373(21):2005-14. Epub 2015/09/29. PMID: 26412349
32. Kalinsky K, Barlow WE, Gralow JR, et al. 21-Gene Assay to Inform Chemotherapy Benefit in Node-Positive Breast Cancer. *N Engl J Med.* 2021;385(25):2336-47. Epub 2021/12/17. PMID: 34914339
33. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016;375(8):717-29. Epub 2016/08/25. PMID: 27557300
34. Piccart M, van 't Veer LJ, Poncet C, et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of the phase 3 randomised MINDACT trial with an exploratory analysis by age. *Lancet Oncol.* 2021;22(4):476-88. Epub 2021/03/16. PMID: 33721561
35. Andre F, Ismaila N, Allison KH, et al. Biomarkers for Adjuvant Endocrine and Chemotherapy in Early-Stage Breast Cancer: ASCO Guideline Update. *J Clin Oncol.* 2022;40(16):1816-37. Epub 2022/04/20. PMID: 35439025
36. Bartlett JMS, Sgroi DC, Treuner K, et al. Breast Cancer Index Is a Predictive Biomarker of Treatment Benefit and Outcome from Extended Tamoxifen Therapy: Final Analysis of the Trans-aTTom Study. *Clin Cancer Res.* 2022;28(9):1871-80. Epub 2022/02/12. PMID: 35144966

37. Sgroi DC, Sestak I, Cuzick J, et al. Prediction of late distant recurrence in patients with oestrogen-receptor-positive breast cancer: a prospective comparison of the breast-cancer index (BCI) assay, 21-gene recurrence score, and IHC4 in the TransATAC study population. *Lancet Oncol.* 2013;14(11):1067-76. Epub 2013/09/17. PMID: 24035531
38. Mosele F, Remon J, Mateo J, et al. Recommendations for the use of next-generation sequencing (NGS) for patients with metastatic cancers: a report from the ESMO Precision Medicine Working Group. *Ann Oncol.* 2020;31(11):1491-505. Epub 2020/08/28. PMID: 32853681
39. Condorelli R, Mosele F, Verret B, et al. Genomic alterations in breast cancer: level of evidence for actionability according to ESMO Scale for Clinical Actionability of molecular Targets (ESCAT). *Ann Oncol.* 2019;30(3):365-73. Epub 2019/02/05. PMID: 30715161
40. André F, Ciruelos E, Rubovszky G, et al. Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med.* 2019;380(20):1929-40. Epub 2019/05/16. PMID: 31091374
41. Robson M, Im SA, Senkus E, et al. Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation. *N Engl J Med.* 2017;377(6):523-33. Epub 2017/06/06. PMID: 28578601
42. Fribbens C, O'Leary B, Kilburn L, et al. Plasma ESR1 Mutations and the Treatment of Estrogen Receptor-Positive Advanced Breast Cancer. *J Clin Oncol.* 2016;34(25):2961-8. Epub 2016/06/09. PMID: 27269946
43. Valle JW, Kelley RK, Nervi B, et al. Biliary tract cancer. *Lancet.* 2021;397(10272):428-44. Epub 2021/02/01. PMID: 33516341
44. Lamarca A, Edeline J, Goyal L. How I treat biliary tract cancer. *ESMO open.* 2022;7(1):100378. Epub 2022/01/16. PMID: 35032765
45. Abou-Alfa GK, Macarulla T, Javle MM, et al. Ivosidenib in IDH1-mutant, chemotherapy-refractory cholangiocarcinoma (ClarIDHy): a multicentre, randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Oncol.* 2020;21(6):796-807. Epub 2020/05/18. PMID: 32416072
46. Biller LH, Schrag D. Diagnosis and Treatment of Metastatic Colorectal Cancer: A Review. *Jama.* 2021;325(7):669-85. Epub 2021/02/17. PMID: 33591350
47. Stadler ZK, Battaglin F, Middha S, et al. Reliable Detection of Mismatch Repair Deficiency in Colorectal Cancers Using Mutational Load in Next-Generation Sequencing Panels. *J Clin Oncol.* 2016;34(18):2141-7. Epub 2016/03/30. PMID: 27022117
48. Lynch HT, Lynch PM. Molecular screening for the Lynch syndrome--better than family history? *N Engl J Med.* 2005;352(18):1920-2. Epub 2005/05/06. PMID: 15872208
49. Ribic CM, Sargent DJ, Moore MJ, et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N Engl J Med.* 2003;349(3):247-57. Epub 2003/07/18. PMID: 12867608
50. Allegra CJ, Rumble RB, Hamilton SR, et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J Clin Oncol.* 2016;34(2):179-85. Epub 2015/10/07. PMID: 26438111
51. Korphaisarn K, Kopetz S. BRAF-Directed Therapy in Metastatic Colorectal Cancer. *Cancer J.* 2016;22(3):175-8. Epub 2016/06/25. PMID: 27341594
52. Corcoran RB, Atreya CE, Falchook GS, et al. Combined BRAF and MEK Inhibition With Dabrafenib and Trametinib in BRAF V600-Mutant Colorectal Cancer. *J Clin Oncol.* 2015;33(34):4023-31. Epub 2015/09/24. PMID: 26392102
53. Tabernero J, Grothey A, Van Cutsem E, et al. Encorafenib Plus Cetuximab as a New Standard of Care for Previously Treated BRAF V600E-Mutant Metastatic Colorectal Cancer: Updated Survival Results and Subgroup Analyses from the BEACON Study. *J Clin Oncol.* 2021;39(4):273-84. Epub 2021/01/28. PMID: 33503393
54. Xu Y, Xing L, Su J, et al. Model-based clustering for identifying disease-associated SNPs in case-control genome-wide association studies. *Sci Rep.* 2019;9(1):13686. Epub 2019/09/25. PMID: 31548641
55. Solomon JP, Hechtman JF. Detection of NTRK Fusions: Merits and Limitations of Current Diagnostic Platforms. *Cancer Res.* 2019;79(13):3163-8. Epub 2019/06/15. PMID: 31196931
56. Miedema J, Andea AA. Through the looking glass and what you find there: making sense of comparative genomic hybridization and fluorescence in situ hybridization for melanoma diagnosis. *Mod Pathol.* 2020;33(7):1318-30. Epub 2020/02/19. PMID: 32066861
57. Clarke LE, Warf MB, Flake DD, 2nd, et al. Clinical validation of a gene expression signature that differentiates benign nevi from malignant melanoma. *J Cutan Pathol.* 2015;42(4):244-52. Epub 2015/03/03. PMID: 25727210
58. Clarke LE, Flake DD, 2nd, Busam K, et al. An independent validation of a gene expression signature to differentiate malignant melanoma from benign melanocytic nevi. *Cancer.* 2017;123(4):617-28. Epub 2016/10/22. PMID: 27768230
59. Ko JS, Clarke LE, Minca EC, et al. Correlation of melanoma gene expression score with clinical outcomes on a series of melanocytic lesions. *Hum Pathol.* 2019;86:213-21. Epub 2018/12/20. PMID: 30566894
60. Chan WH, Tsao H. Consensus, Controversy, and Conversations About Gene Expression Profiling in Melanoma. *JAMA Dermatol.* 2020;156(9):949-51. Epub 2020/08/04. PMID: 32745178
61. Greenhaw BN, Covington KR, Kurley SJ, et al. Molecular risk prediction in cutaneous melanoma: A meta-analysis of the 31-gene expression profile prognostic test in 1,479 patients. *J Am Acad Dermatol.* 2020;83(3):745-53. Epub 2020/04/02. PMID: 32229276

62. Marchetti MA, Dusza SW, Bartlett EK. Problematic methodology in a systematic review and meta-analysis of DecisionDx-Melanoma. *J Am Acad Dermatol*. 2020;83(5):e357-e8. Epub 2020/06/12. PMID: 32526321
63. Kovarik CL, Chu EY, Adamson AS. Gene Expression Profile Testing for Thin Melanoma: Evidence to Support Clinical Use Remains Thin. *JAMA Dermatol*. 2020;156(8):837-8. Epub 2020/04/16. PMID: 32293654
64. Seider MI, Mruthyunjaya P. MOLECULAR PROGNOSTICS FOR UVEAL MELANOMA. *Retina*. 2018;38(2):211-9. Epub 2017/07/01. PMID: 28665870
65. Chattopadhyay C, Kim DW, Gombos DS, et al. Uveal melanoma: From diagnosis to treatment and the science in between. *Cancer*. 2016;122(15):2299-312. Epub 2016/03/19. PMID: 26991400
66. Aaberg TM, Covington KR, Tsai T, et al. Gene Expression Profiling in Uveal Melanoma: Five-Year Prospective Outcomes and Meta-Analysis. *Ocul Oncol Pathol*. 2020;6(5):360-7. Epub 2020/10/31. PMID: 33123530
67. Francis JH, Patel SP, Gombos DS, et al. Surveillance options for patients with uveal melanoma following definitive management. *Am Soc Clin Oncol Educ Book*. 2013:382-7. Epub 2013/05/30. PMID: 23714555
68. Cheng L, Lopez-Beltran A, Massari F, et al. Molecular testing for BRAF mutations to inform melanoma treatment decisions: a move toward precision medicine. *Mod Pathol*. 2018;31(1):24-38. Epub 2017/11/18. PMID: 29148538
69. Michielin O, van Akkooi ACJ, Ascierto PA, et al. Cutaneous melanoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up†. *Ann Oncol*. 2019;30(12):1884-901. Epub 2019/10/01. PMID: 31566661
70. Nathan P, Hassel JC, Rutkowski P, et al. Overall Survival Benefit with Tebentafusp in Metastatic Uveal Melanoma. *N Engl J Med*. 2021;385(13):1196-206. Epub 2021/09/23. PMID: 34551229
71. Devitt B, Liu W, Salemi R, et al. Clinical outcome and pathological features associated with NRAS mutation in cutaneous melanoma. *Pigment Cell Melanoma Res*. 2011;24(4):666-72. Epub 2011/05/28. PMID: 21615881
72. Dummer R, Schadendorf D, Ascierto PA, et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2017;18(4):435-45. Epub 2017/03/13. PMID: 28284557
73. Steeb T, Wessely A, Petzold A, et al. c-Kit inhibitors for unresectable or metastatic mucosal, acral or chronically sun-damaged melanoma: a systematic review and one-arm meta-analysis. *Eur J Cancer*. 2021;157:348-57. Epub 2021/09/26. PMID: 34562816
74. Griesinger F, Eberhardt W, Nusch A, et al. Biomarker testing in non-small cell lung cancer in routine care: Analysis of the first 3,717 patients in the German prospective, observational, nation-wide CRISP Registry (AIO-TRK-0315). *Lung Cancer*. 2021;152:174-84. Epub 2020/12/29. PMID: 33358484
75. Tan AC, Tan DSW. Targeted Therapies for Lung Cancer Patients With Oncogenic Driver Molecular Alterations. *J Clin Oncol*. 2022;40(6):611-25. Epub 2022/01/06. PMID: 34985916
76. Kalemkerian GP, Narula N, Kennedy EB, et al. Molecular Testing Guideline for the Selection of Patients With Lung Cancer for Treatment With Targeted Tyrosine Kinase Inhibitors: American Society of Clinical Oncology Endorsement of the College of American Pathologists/International Association for the Study of Lung Cancer/Association for Molecular Pathology Clinical Practice Guideline Update. *J Clin Oncol*. 2018;36(9):911-9. Epub 2018/02/06. PMID: 29401004
77. Ionescu DN, Stockley TL, Banerji S, et al. Consensus Recommendations to Optimize Testing for New Targetable Alterations in Non-Small Cell Lung Cancer. *Curr Oncol*. 2022;29(7):4981-97. Epub 2022/07/26. PMID: 35877256
78. Wu YL, Tsuboi M, He J, et al. Osimertinib in Resected EGFR-Mutated Non-Small-Cell Lung Cancer. *N Engl J Med*. 2020;383(18):1711-23. Epub 2020/09/22. PMID: 32955177
79. Sotelo MJ, Luis García J, Torres-Mattos C, et al. Recent advances and new insights in the management of early-stage epidermal growth factor receptor-mutated non-small-cell lung cancer. *World J Clin Oncol*. 2021;12(10):912-25. Epub 2021/11/05. PMID: 34733613
80. Singhi EK, Gay CM. Narrative review of the emerging role of molecular biomarkers in guiding the definitive management of unresectable non-small cell lung cancer. *Transl Lung Cancer Res*. 2020;9(5):2051-8. Epub 2020/11/20. PMID: 33209625
81. Dizon DS. PARP inhibitors for targeted treatment in ovarian cancer. *Lancet*. 2017;390(10106):1929-30. Epub 2017/09/17. PMID: 28916370
82. Ledermann JA, Oza AM, Lorusso D, et al. Rucaparib for patients with platinum-sensitive, recurrent ovarian carcinoma (ARIEL3): post-progression outcomes and updated safety results from a randomised, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2020;21(5):710-22. Epub 2020/05/04. PMID: 32359490
83. Coleman RL, Oza AM, Lorusso D, et al. Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2017;390(10106):1949-61. Epub 2017/09/17. PMID: 28916367
84. Rainone M, Singh I, Salo-Mullen EE, et al. An Emerging Paradigm for Germline Testing in Pancreatic Ductal Adenocarcinoma and Immediate Implications for Clinical Practice: A Review. *JAMA Oncol*. 2020;6(5):764-71. Epub 2020/02/14. PMID: 32053139
85. Golan T, Hammel P. Management of BRCA Mutation Carriers With Pancreatic Adenocarcinoma. *J Natl Compr Canc Netw*. 2021;19(4):469-73. Epub 2021/04/13. PMID: 33845459
86. Park W, Chawla A, O'Reilly EM. Pancreatic Cancer: A Review. *Jama*. 2021;326(9):851-62. Epub 2021/09/22. PMID: 34547082

87. Golan T, Hammel P, Reni M, et al. Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer. *N Engl J Med*. 2019;381(4):317-27. Epub 2019/06/04. PMID: 31157963
88. Kindler HL, Hammel P, Reni M, et al. Overall Survival Results From the POLO Trial: A Phase III Study of Active Maintenance Olaparib Versus Placebo for Germline BRCA-Mutated Metastatic Pancreatic Cancer. *J Clin Oncol*. 2022;Jco2101604. Epub 2022/07/15. PMID: 35834777
89. Nishikawa G, Booth C, Prasad V. Olaparib for BRCA mutant pancreas cancer: Should the POLO trial change clinical practice? *Cancer*. 2020;126(18):4087-8. Epub 2020/05/28. PMID: 32459369
90. Reiss KA, Mick R, O'Hara MH, et al. Phase II Study of Maintenance Rucaparib in Patients With Platinum-Sensitive Advanced Pancreatic Cancer and a Pathogenic Germline or Somatic Variant in BRCA1, BRCA2, or PALB2. *J Clin Oncol*. 2021;39(22):2497-505. Epub 2021/05/11. PMID: 33970687
91. Mottet N, Bellmunt J, Bolla M, et al. EAU-ESTRO-SIOG Guidelines on Prostate Cancer. Part 1: Screening, Diagnosis, and Local Treatment with Curative Intent. *Eur Urol*. 2017;71(4):618-29. Epub 2016/08/30. PMID: 27568654
92. Eggener SE, Berlin A, Vickers AJ, et al. Low-Grade Prostate Cancer: Time to Stop Calling It Cancer. *J Clin Oncol*. 2022;Jco2200123. Epub 2022/04/19. PMID: 35436144
93. Zelic R, Garmo H, Zugna D, et al. Predicting Prostate Cancer Death with Different Pretreatment Risk Stratification Tools: A Head-to-head Comparison in a Nationwide Cohort Study. *Eur Urol*. 2020;77(2):180-8. Epub 2019/10/14. PMID: 31606332
94. Haffner MC, Zwart W, Roudier MP, et al. Genomic and phenotypic heterogeneity in prostate cancer. *Nat Rev Urol*. 2021;18(2):79-92. Epub 2020/12/18. PMID: 33328650
95. Eggener SE, Rumble RB, Armstrong AJ, et al. Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. *J Clin Oncol*. 2020;38(13):1474-94. Epub 2019/12/13. PMID: 31829902
96. National Comprehensive Cancer Network (NCCN). Prostate Cancer Version 4.2022, NCCN Clinical Practice Guidelines in Oncology. 2022.
97. Lin DW, Nelson PS. Prognostic Genomic Biomarkers in Patients With Localized Prostate Cancer: Is Rising Utilization Justified by Evidence? *JAMA Oncol*. 2021;7(1):59-60. Epub 2020/11/26. PMID: 33237305
98. Leapman MS, Wang R, Ma S, et al. Regional Adoption of Commercial Gene Expression Testing for Prostate Cancer. *JAMA Oncol*. 2021;7(1):52-8. Epub 2020/11/26. PMID: 33237277
99. Hu JC, Tosoian JJ, Qi J, et al. Clinical Utility of Gene Expression Classifiers in Men With Newly Diagnosed Prostate Cancer. *JCO Precis Oncol*. 2018;2. Epub 2018/01/01. PMID: 32832833
100. Mateo J, McKay R, Abida W, et al. Accelerating precision medicine in metastatic prostate cancer. *Nat Cancer*. 2020;1(11):1041-53. Epub 2021/07/15. PMID: 34258585
101. Giri VN, Morgan TM, Morris DS, et al. Genetic testing in prostate cancer management: Considerations informing primary care. *CA Cancer J Clin*. 2022;72(4):360-71. Epub 2022 Feb 25. PMID: 35201622
102. Sokol ES, Jin DX, Fine A, et al. PARP Inhibitor Insensitivity to BRCA1/2 Monoallelic Mutations in Microsatellite Instability-High Cancers. *JCO Precis Oncol*. 2022;6:e2100531. Epub 2022/07/01. PMID: 35772050
103. de Bono J, Mateo J, Fizazi K, et al. Olaparib for Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med*. 2020;382(22):2091-102. Epub 2020/04/29. PMID: 32343890
104. Hussain M, Mateo J, Fizazi K, et al. Survival with Olaparib in Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med*. 2020;383(24):2345-57. Epub 2020/09/22. PMID: 32955174
105. Abida W, Patnaik A, Campbell D, et al. Rucaparib in Men With Metastatic Castration-Resistant Prostate Cancer Harboring a BRCA1 or BRCA2 Gene Alteration. *J Clin Oncol*. 2020;38(32):3763-72. Epub 2020/08/17. PMID: 32795228
106. Abida W, Campbell D, Patnaik A, et al. Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study. *Clin Cancer Res*. 2020;26(11):2487-96. Epub 2020/02/23. PMID: 32086346
107. Hanba C, Khariwala SS. What is the Utility of Genetic Testing in Indeterminate Thyroid Nodules? *The Laryngoscope*. 2021;131(11):2399-400. Epub 2021/02/10. PMID: 33559898
108. Leboulleux S, Bournaud C, Chougnet CN, et al. Thyroidectomy without Radioiodine in Patients with Low-Risk Thyroid Cancer. *N Engl J Med*. 2022;386(10):923-32. Epub 2022/03/10. PMID: 35263518
109. Zanocco KA, Hershman JM, Leung AM. Active Surveillance of Low-Risk Thyroid Cancer. *Jama*. 2019;321(20):2020-1. Epub 2019/05/01. PMID: 31038662
110. Cibas ES, Ali SZ. The 2017 Bethesda System for Reporting Thyroid Cytopathology. *Thyroid*. 2017;27(11):1341-6. Epub 2017/11/02. PMID: 29091573
111. Deaver KE, Haugen BR, Pozdeyev N, et al. Outcomes of Bethesda categories III and IV thyroid nodules over 5 years and performance of the Afirma gene expression classifier: A single-institution study. *Clin Endocrinol (Oxf)*. 2018;89(2):226-32. Epub 2018/05/24. PMID: 29791966
112. Mayson SE, Haugen BR. Molecular Diagnostic Evaluation of Thyroid Nodules. *Endocrinol Metab Clin North Am*. 2019;48(1):85-97. Epub 2019/02/06. PMID: 30717912

113. Sistrunk JW, Shifrin A, Frager M, et al. Clinical impact of testing for mutations and microRNAs in thyroid nodules. *Diagn Cytopathol.* 2019;47(8):758-64. Epub 2019/04/24. PMID: 31013001
114. Lupo MA, Walts AE, Sistrunk JW, et al. Multiplatform molecular test performance in indeterminate thyroid nodules. *Diagn Cytopathol.* 2020;48(12):1254-64. Epub 2020/08/09. PMID: 32767735
115. Angell TE, Wirth LJ, Cabanillas ME, et al. Analytical and Clinical Validation of Expressed Variants and Fusions From the Whole Transcriptome of Thyroid FNA Samples. *Front Endocrinol (Lausanne).* 2019;10:612. Epub 2019/10/02. PMID: 31572297
116. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid.* 2016;26(1):1-133. Epub 2015/10/16. PMID: 26462967
117. Marti JL, Avadhani V, Donatelli LA, et al. Wide Inter-institutional Variation in Performance of a Molecular Classifier for Indeterminate Thyroid Nodules. *Ann Surg Oncol.* 2015;22(12):3996-4001. Epub 2015/04/12. PMID: 25862581
118. Luster M, Aktolun C, Amendoeira I, et al. European Perspective on 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: Proceedings of an Interactive International Symposium. *Thyroid.* 2019;29(1):7-26. Epub 2018/11/30. PMID: 30484394
119. Twining CL, Lupo MA, Tuttle RM. Implementing Key Changes in the American Thyroid Association 2015 Thyroid Nodules/Differentiated Thyroid Cancer Guidelines Across Practice Types. *Endocr Pract.* 2018;24(9):833-40. Epub 2018/10/12. PMID: 30308136
120. Steward DL, Carty SE, Sippel RS, et al. Performance of a Multigene Genomic Classifier in Thyroid Nodules With Indeterminate Cytology: A Prospective Blinded Multicenter Study. *JAMA Oncol.* 2019;5(2):204-12. Epub 2018/11/13. PMID: 30419129
121. Valderrabano P, Hallanger-Johnson JE, Thapa R, et al. Comparison of Postmarketing Findings vs the Initial Clinical Validation Findings of a Thyroid Nodule Gene Expression Classifier: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg.* 2019;145(9):783-92. Epub 2019/07/19. PMID: 31318389
122. Hayes DF. Defining Clinical Utility of Tumor Biomarker Tests: A Clinician's Viewpoint. *J Clin Oncol.* 2021;39(3):238-48. Epub 2020 Dec 17. PMID: 33326253
123. Marti JL, Shaha AR. Molecular Testing for Indeterminate Thyroid Nodules-When the Rubber Meets the Road. *JAMA Otolaryngol Head Neck Surg.* 2019;145(9):792-3. Epub 2019/07/19. PMID: 31318432
124. Doerfler WR, Nikitski AV, Morariu EM, et al. Molecular alterations in Hürthle cell nodules and preoperative cancer risk. *Endocr Relat Cancer.* 2021;28(5):301-9. Epub 2021/04/02. PMID: 33792557
125. Schatz-Siemers N, Brandler TC, Oweity T, et al. Hürthle cell lesions on thyroid fine needle aspiration cytology: Molecular and histologic correlation. *Diagn Cytopathol.* 2019;47(10):977-85. Epub 2019/07/12. PMID: 31293091
126. Hao Y, Duh QY, Kloos RT, et al. Identification of Hürthle cell cancers: solving a clinical challenge with genomic sequencing and a trio of machine learning algorithms. *BMC Syst Biol.* 2019;13(Suppl 2):27. Epub 2019/04/07. PMID: 30952205
127. Rassy E, Pavlidis N. The currently declining incidence of cancer of unknown primary. *Cancer Epidemiol.* 2019;61:139-41. Epub 2019/06/30. PMID: 31254795
128. Rassy E, Assi T, Pavlidis N. Exploring the biological hallmarks of cancer of unknown primary: where do we stand today? *Br J Cancer.* 2020;122(8):1124-32. Epub 2020/02/12. PMID: 32042068
129. Varadhachary GR, Raber MN. Cancer of unknown primary site. *N Engl J Med.* 2014;371(8):757-65. Epub 2014/08/21. PMID: 25140961
130. Huey RW, Smaglo BG, Estrella JS, et al. Cancer of Unknown Primary Presenting as Bone-Predominant or Lymph Node-Only Disease: A Clinicopathologic Portrait. *Oncologist.* 2021;26(4):e650-e7. Epub 2021/02/02. PMID: 33524217
131. Naing A, Meric-Bernstam F, Stephen B, et al. Phase 2 study of pembrolizumab in patients with advanced rare cancers. *J Immunother Cancer.* 2020;8(1). Epub 2020/03/20. PMID: 32188704
132. Nguyen L, Van Hoeck A, Cuppen E. Machine learning-based tissue of origin classification for cancer of unknown primary diagnostics using genome-wide mutation features. *Nat Commun.* 2022;13(1):4013. Epub 2022/07/12. PMID: 35817764
133. Jiao W, Atwal G, Polak P, et al. A deep learning system accurately classifies primary and metastatic cancers using passenger mutation patterns. *Nat Commun.* 2020;11(1):728. Epub 2020/02/07. PMID: 32024849
134. Salvadores M, Mas-Ponte D, Supek F. Passenger mutations accurately classify human tumors. *PLoS Comput Biol.* 2019;15(4):e1006953. Epub 2019/04/16. PMID: 30986244
135. Raghav K, Hwang H, Jácome AA, et al. Development and Validation of a Novel Nomogram for Individualized Prediction of Survival in Cancer of Unknown Primary. *Clin Cancer Res.* 2021;27(12):3414-21. Epub 2021/04/17. PMID: 33858857
136. Hunger SP, Mullighan CG. Acute Lymphoblastic Leukemia in Children. *N Engl J Med.* 2015;373(16):1541-52. Epub 2015/10/16. PMID: 26465987
137. Jabbour E, Pui CH, Kantarjian H. Progress and Innovations in the Management of Adult Acute Lymphoblastic Leukemia. *JAMA Oncol.* 2018;4(10):1413-20. Epub 2018/06/23. PMID: 29931220
138. Liu YF, Wang BY, Zhang WN, et al. Genomic Profiling of Adult and Pediatric B-cell Acute Lymphoblastic Leukemia. *EBioMedicine.* 2016;8:173-83. Epub 2016/07/20. PMID: 27428428

139. Berry DA, Zhou S, Higley H, et al. Association of Minimal Residual Disease With Clinical Outcome in Pediatric and Adult Acute Lymphoblastic Leukemia: A Meta-analysis. *JAMA Oncol.* 2017;3(7):e170580. Epub 2017/05/12. PMID: 28494052
140. Vora A, Goulden N, Mitchell C, et al. Augmented post-remission therapy for a minimal residual disease-defined high-risk subgroup of children and young people with clinical standard-risk and intermediate-risk acute lymphoblastic leukaemia (UKALL 2003): a randomised controlled trial. *Lancet Oncol.* 2014;15(8):809-18. Epub 2014/06/14. PMID: 24924991
141. Brown P, Inaba H, Annesley C, et al. Pediatric Acute Lymphoblastic Leukemia, Version 2.2020, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw.* 2020;18(1):81-112. Epub 2020/01/08. PMID: 31910389
142. Aitken MJL, Ravandi F, Patel KP, et al. Prognostic and therapeutic implications of measurable residual disease in acute myeloid leukemia. *J Hematol Oncol.* 2021;14(1):137. Epub 2021/09/05. PMID: 34479626
143. Döhner H, Weisdorf DJ, Bloomfield CD. Acute Myeloid Leukemia. *N Engl J Med.* 2015;373(12):1136-52. Epub 2015/09/17. PMID: 26376137
144. Nabhan C, Kamat S, Karl Kish J. Acute myeloid leukemia in the elderly: what constitutes treatment value? *Leuk Lymphoma.* 2019;60(5):1164-70. Epub 2018/11/09. PMID: 30407103
145. Khoury JD, Solary E, Abla O, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia.* 2022;36(7):1703-19. Epub 2022/06/23. PMID: 35732831
146. Duncavage EJ, Schroeder MC, O'Laughlin M, et al. Genome Sequencing as an Alternative to Cytogenetic Analysis in Myeloid Cancers. *N Engl J Med.* 2021;384(10):924-35. Epub 2021/03/12. PMID: 33704937
147. Döhner H, Wei AH, Appelbaum FR, et al. Diagnosis and Management of AML in Adults: 2022 ELN Recommendations from an International Expert Panel. *Blood.* 2022. Epub 2022/07/08. PMID: 35797463
148. Short NJ, Tallman MS, Pollyea DA, et al. Optimizing Risk Stratification in Acute Myeloid Leukemia: Dynamic Models for a Dynamic Therapeutic Landscape. *J Clin Oncol.* 2021;39(23):2535-8. Epub 2021/05/28. PMID: 34043455
149. Ivey A, Hills RK, Simpson MA, et al. Assessment of Minimal Residual Disease in Standard-Risk AML. *N Engl J Med.* 2016;374(5):422-33. Epub 2016/01/21. PMID: 26789727
150. Short NJ, Zhou S, Fu C, et al. Association of Measurable Residual Disease With Survival Outcomes in Patients With Acute Myeloid Leukemia: A Systematic Review and Meta-analysis. *JAMA Oncol.* 2020;6(12):1890-9. Epub 2020/10/09. PMID: 33030517
151. Dillon R, Potter N, Freeman S, et al. How we use molecular minimal residual disease (MRD) testing in acute myeloid leukaemia (AML). *Br J Haematol.* 2021;193(2):231-44. Epub 2020/10/16. PMID: 33058194
152. Spivak JL. Myeloproliferative Neoplasms. *N Engl J Med.* 2017;377(9):895-6. Epub 2017/08/31. PMID: 28854086
153. Hochhaus A, Saussele S, Rosti G, et al. Chronic myeloid leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2018;29(Suppl 4):iv261. Epub 2018/10/05. PMID: 30285223
154. Wong WJ, Pozdnyakova O. Myeloproliferative neoplasms: Diagnostic workup of the cythemic patient. *Int J Lab Hematol.* 2019;41 Suppl 1:142-50. Epub 2019/05/10. PMID: 31069979
155. Mughal TI, Gotlib J, Mesa R, et al. Recent advances in the genomics and therapy of BCR/ABL1-positive and -negative chronic myeloproliferative neoplasms. *Leuk Res.* 2018;67:67-74. Epub 2018/02/22. PMID: 29466766
156. Hochhaus A, Larson RA, Guilhot F, et al. Long-Term Outcomes of Imatinib Treatment for Chronic Myeloid Leukemia. *N Engl J Med.* 2017;376(10):917-27. Epub 2017/03/09. PMID: 28273028
157. Cross NC, White HE, Colomer D, et al. Laboratory recommendations for scoring deep molecular responses following treatment for chronic myeloid leukemia. *Leukemia.* 2015;29(5):999-1003. Epub 2015/02/06. PMID: 25652737
158. Shanmuganathan N, Hughes TP. Molecular monitoring in CML: how deep? How often? How should it influence therapy? *Hematology Am Soc Hematol Educ Program.* 2018;2018(1):168-76. Epub 2018/12/07. PMID: 30504306
159. Hochhaus A, Baccarani M, Silver RT, et al. European LeukemiaNet 2020 recommendations for treating chronic myeloid leukemia. *Leukemia.* 2020;34(4):966-84. Epub 2020/03/05. PMID: 32127639
160. Soverini S, Bavaro L, De Benedittis C, et al. Prospective assessment of NGS-detectable mutations in CML patients with nonoptimal response: the NEXT-in-CML study. *Blood.* 2020;135(8):534-41. Epub 2019/12/27. PMID: 31877211
161. Rajkumar SV. Multiple myeloma: 2020 update on diagnosis, risk-stratification and management. *Am J Hematol.* 2020;95(5):548-67. Epub 2020/03/27. PMID: 32212178
162. Mikhael J, Ismaila N, Cheung MC, et al. Treatment of Multiple Myeloma: ASCO and CCO Joint Clinical Practice Guideline. *J Clin Oncol.* 2019;37(14):1228-63. Epub 2019/04/02. PMID: 30932732
163. Tandon N, Rajkumar SV, LaPlant B, et al. Clinical utility of the Revised International Staging System in unselected patients with newly diagnosed and relapsed multiple myeloma. *Blood Cancer J.* 2017;7(2):e528. Epub 2017/02/18. PMID: 28211889
164. Abdallah NH, Binder M, Rajkumar SV, et al. A simple additive staging system for newly diagnosed multiple myeloma. *Blood Cancer J.* 2022;12(1):21. Epub 2022/02/02. PMID: 35102148
165. Kumar S, Paiva B, Anderson KC, et al. International Myeloma Working Group consensus criteria for response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol.* 2016;17(8):e328-e46. Epub 2016/08/12. PMID: 27511158

166. Dimopoulos MA, Moreau P, Terpos E, et al. Multiple myeloma: EHA-ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up(+). *Ann Oncol.* 2021;32(3):309-22. Epub 2021/02/08. PMID: 33549387
167. Cazzola M. Myelodysplastic Syndromes. *N Engl J Med.* 2020;383(14):1358-74. Epub 2020/10/01. PMID: 32997910
168. Fenaux P, Platzbecker U, Ades L. How we manage adults with myelodysplastic syndrome. *Br J Haematol.* 2020;189(6):1016-27. Epub 2019/10/01. PMID: 31568568
169. National Comprehensive Cancer Network (NCCN). NCCN Guidelines®: Myelodysplastic Syndromes, Version 3.2022. 2022.
170. Garcia-Manero G, Chien KS, Montalban-Bravo G. Myelodysplastic syndromes: 2021 update on diagnosis, risk stratification and management. *Am J Hematol.* 2020;95(11):1399-420. Epub 2020/08/04. PMID: 32744763
171. Bernard E, Tuechler H, Greenberg Peter L, et al. Molecular International Prognostic Scoring System for Myelodysplastic Syndromes. *NEJM Evid.* 2022;1(7):EVIDoa2200008.

Codes

The following code list is not meant to be all-inclusive. Authorization requirements will vary by health plan. Please consult the applicable health plan for guidance on specific procedure codes.

Specific CPT codes for services should be used when available. Nonspecific or not otherwise classified codes may be subject to additional documentation requirements and review.

CPT/HCPCS

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright by the American Medical Association. All Rights Reserved. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

81120	IDH1 (isocitrate dehydrogenase 1 [NADP+], soluble) (eg, glioma), common variants (eg, R132H, R132C)
81121	IDH2 (isocitrate dehydrogenase 2 [NADP+], mitochondrial) (eg, glioma), common variants (eg, R140W, R172M)
81162	BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full sequence analysis and full duplication/deletion analysis (ie, detection of large gene rearrangements)
81163	BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full sequence analysis
81164	BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full duplication/deletion analysis (ie, detection of large gene rearrangements)
81168	CCND1/IGH (t(11;14)) (eg, mantle cell lymphoma) translocation analysis, major breakpoint, qualitative and quantitative, if performed
81170	ABL1 (ABL proto-oncogene 1, non-receptor tyrosine kinase) (eg, acquired imatinib tyrosine kinase inhibitor resistance), gene analysis, variants in the kinase domain
81173	AR (androgen receptor) (eg, spinal and bulbar muscular atrophy, Kennedy disease, X chromosome inactivation) gene analysis; full gene sequence
81175	ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; full gene sequence
81176	ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; targeted sequence analysis (eg, exon 12)
81191	NTRK1 (neurotrophic receptor tyrosine kinase 1) (eg, solid tumors) translocation analysis
81192	NTRK2 (neurotrophic receptor tyrosine kinase 2) (eg, solid tumors) translocation analysis
81193	NTRK3 (neurotrophic receptor tyrosine kinase 3) (eg, solid tumors) translocation analysis
81194	NTRK (neurotrophic-tropomyosin receptor tyrosine kinase 1, 2, and 3) (eg, solid tumors) translocation analysis
81206	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; major breakpoint, qualitative or quantitative
81207	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; minor breakpoint, qualitative or quantitative
81208	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; other breakpoint, qualitative or quantitative
81210	BRAF (B-Raf proto-oncogene, serine/threonine kinase) (eg, colon cancer, melanoma), gene analysis, V600 variant(s)
81218	CEBPA (CCAAT/enhancer binding protein [C/EBP], alpha) (eg, acute myeloid leukemia), gene analysis, full gene sequence
81219	CALR (calreticulin) (eg, myeloproliferative disorders), gene analysis, common variants in exon 9
81233	BTK (Bruton's tyrosine kinase) (eg, chronic lymphocytic leukemia) gene analysis, common variants (eg, C481S, C481R, C481F)

81235	EGFR (epidermal growth factor receptor) (eg, non-small cell lung cancer) gene analysis, common variants (eg, exon 19 LREA deletion, L858R, T790M, G719A, G719S, L861Q)
81236	EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit) (eg, myelodysplastic syndrome, myeloproliferative neoplasms) gene analysis, full gene sequence
81237	EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit) (eg, diffuse large B-cell lymphoma) gene analysis, common variant(s) (eg, codon 646)
81245	FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; internal tandem duplication (ITD) variants (ie, exons 14, 15)
81246	FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; tyrosine kinase domain (TKD) variants (eg, D835, I836)
81261	IGH@ (Immunoglobulin heavy chain locus) (eg, leukemias and lymphomas, B-cell), gene rearrangement analysis to detect abnormal clonal population(s); amplified methodology (eg, polymerase chain reaction)
81262	IGH@ (Immunoglobulin heavy chain locus) (eg, leukemias and lymphomas, B-cell), gene rearrangement analysis to detect abnormal clonal population(s); direct probe methodology (eg, Southern blot)
81263	IGH@ (Immunoglobulin heavy chain locus) (eg, leukemia and lymphoma, B-cell), variable region somatic mutation analysis
81264	IGK@ (Immunoglobulin kappa light chain locus) (eg, leukemia and lymphoma, B-cell), gene rearrangement analysis, evaluation to detect abnormal clonal population(s)
81270	JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) gene analysis, p.Val617Phe (V617F) variant
81272	KIT (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, gastrointestinal stromal tumor [GIST], acute myeloid leukemia, melanoma), gene analysis, targeted sequence analysis (eg, exons 8, 11, 13, 17, 18)
81273	KIT (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, mastocytosis), gene analysis, D816 variant(s)
81275	KRAS (Kirsten rat sarcoma viral oncogene homolog) (eg, carcinoma) gene analysis; variants in exon 2 (eg, codons 12 and 13)
81276	KRAS (Kirsten rat sarcoma viral oncogene homolog) (eg, carcinoma) gene analysis; additional variant(s) (eg, codon 61, codon 146)
81277	Cytogenomic neoplasia (genome-wide) microarray analysis, interrogation of genomic regions for copy number and loss-of-heterozygosity variants for chromosomal abnormalities
81278	IGH@/BCL2 (t(14;18)) (eg, follicular lymphoma) translocation analysis, major breakpoint region (MBR) and minor cluster region (mcr) breakpoints, qualitative or quantitative
81279	JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) targeted sequence analysis (eg, exons 12 and 13)
81287	MGMT (O-6-methylguanine-DNA methyltransferase) (eg, glioblastoma multiforme) promoter methylation analysis
81288	MLH1 (mutL homolog 1, colon cancer, nonpolyposis type 2) (eg, hereditary non-polyposis colorectal cancer, Lynch syndrome) gene analysis; promoter methylation analysis
81301	Microsatellite instability analysis (eg, hereditary non-polyposis colorectal cancer, Lynch syndrome) of markers for mismatch repair deficiency (eg, BAT25, BAT26), includes comparison of neoplastic and normal tissue, if performed
81305	MYD88 (myeloid differentiation primary response 88) (eg, Waldenstrom's macroglobulinemia, lymphoplasmacytic leukemia) gene analysis, p.Leu265Pro (L265P) variant
81307	<i>PALB2</i> (<i>partner and localizer of BRCA2</i>) (eg, breast and pancreatic cancer) gene analysis; full gene sequence
81309	PIK3CA (phosphatidylinositol-4, 5-biphosphate 3-kinase, catalytic subunit alpha) (eg, colorectal and breast cancer) gene analysis, targeted sequence analysis (eg, exons 7, 9, 20)
81310	NPM1 (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, exon 12 variants
81311	NRAS (neuroblastoma RAS viral [v-ras] oncogene homolog) (eg, colorectal carcinoma), gene analysis, variants in exon 2 (eg, codons 12 and 13) and exon 3 (eg, codon 61)
81313	PCA3/KLK3 (prostate cancer antigen 3 [non-protein coding]/kallikrein-related peptidase 3 [prostate specific antigen]) ratio (eg, prostate cancer)
81314	PDGFRA (platelet-derived growth factor receptor, alpha polypeptide) (eg, gastrointestinal stromal tumor [GIST]), gene analysis, targeted sequence analysis (eg, exons 12, 18)
81315	PML/RARalpha, (t(15;17)), (promyelocytic leukemia/retinoic acid receptor alpha) (eg, promyelocytic leukemia) translocation analysis; common breakpoints (eg, intron 3 and intron 6), qualitative or quantitative
81316	PML/RARalpha, (t(15;17)), (promyelocytic leukemia/retinoic acid receptor alpha) (eg, promyelocytic leukemia) translocation analysis; single breakpoint (eg, intron 3, intron 6 or exon 6), qualitative or quantitative
81320	PLCG2 (phospholipase C gamma 2) (eg, chronic lymphocytic leukemia) gene analysis, common variants (eg, R665W, S707F, L845F)
81321	PTEN (phosphatase and tensin homolog) (eg, Cowden syndrome, PTEN hamartoma tumor syndrome) gene analysis; full sequence analysis
81323	PTEN (phosphatase and tensin homolog) (eg, Cowden syndrome, PTEN hamartoma tumor syndrome) gene analysis; duplication/deletion variant
81327	SEPT9 (Septin9) (eg, colorectal cancer) promoter methylation analysis

81334	RUNX1 (runt related transcription factor 1) (eg, acute myeloid leukemia, familial platelet disorder with associated myeloid malignancy), gene analysis, targeted sequence analysis (eg, exons 3-8)
81338	MPL (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; common variants (eg, W515A, W515K, W515L, W515R)
81339	MPL (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; sequence analysis, exon 10
81340	TRB@ (T cell antigen receptor, beta) (eg, leukemia and lymphoma), gene rearrangement analysis to detect abnormal clonal population(s); using amplification methodology (eg, polymerase chain reaction)
81341	TRB@ (T cell antigen receptor, beta) (eg, leukemia and lymphoma), gene rearrangement analysis to detect abnormal clonal population(s); using direct probe methodology (eg, Southern blot)
81342	TRG@ (T cell antigen receptor, gamma) (eg, leukemia and lymphoma), gene rearrangement analysis, evaluation to detect abnormal clonal population(s)
81345	TERT (telomerase reverse transcriptase) (eg, thyroid carcinoma, glioblastoma multiforme) gene analysis, targeted sequence analysis (eg, promoter region)
81347	SF3B1 (splicing factor [3b] subunit B1) (eg, myelodysplastic syndrome/acute myeloid leukemia) gene analysis, common variants (eg, A672T, E622D, L833F, R625C, R625L)
81348	SRSF2 (serine and arginine-rich splicing factor 2) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variants (eg, P95H, P95L)
81351	TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; full gene sequence
81352	TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; targeted sequence analysis (eg, 4 oncology)
81353	TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; known familial variant
81357	U2AF1 (U2 small nuclear RNA auxiliary factor 1) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variants (eg, S34F, S34Y, Q157R, Q157P)
81360	ZRSR2 (zinc finger CCH-type, RNA binding motif and serine/arginine-rich 2) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variant(s) (eg, E65fs, E122fs, R448fs)
81401	Molecular pathology procedure, Level 2 (eg, 2-10 SNPs, 1 methylated variant, or 1 somatic variant [typically using nonsequencing target variant analysis], or detection of a dynamic mutation disorder/triplet repeat)
81402	Molecular pathology procedure, Level 3 (eg, >10 SNPs, 2-10 methylated variants, or 2-10 somatic variants [typically using non-sequencing target variant analysis], immunoglobulin and T-cell receptor gene rearrangements, duplication/deletion variants of 1 exon, loss of heterozygosity [LOH], uniparental disomy [UPD])
81403	Molecular pathology procedure, Level 4 (eg, analysis of single exon by DNA sequence analysis, analysis of >10 amplicons using multiplex PCR in 2 or more independent reactions, mutation scanning or duplication/deletion variants of 2-5 exons)
81404	Molecular pathology procedure, Level 5 (eg, analysis of 2-5 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 6-10 exons, or characterization of a dynamic mutation disorder/triplet repeat by Southern blot analysis)
81405	Molecular pathology procedure, Level 6 (eg, analysis of 6-10 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 11-25 exons, regionally targeted cytogenomic array analysis)
81406	Molecular pathology procedure, Level 7 (eg, analysis of 11-25 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 26-50 exons, cytogenomic array analysis for neoplasia)
81408	Molecular pathology procedure, Level 9 (eg, analysis of >50 exons in a single gene by DNA sequence analysis)
81445	Targeted genomic sequence analysis panel, solid organ neoplasm, 5-50 genes (eg, ALK, BRAF, CDKN2A, EGFR, ERBB2, KIT, KRAS, MET, NRAS, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed; DNA analysis or combined DNA and RNA analysis
81449	Targeted genomic sequence analysis panel, solid organ neoplasm, 5-50 genes (eg, ALK, BRAF, CDKN2A, EGFR, ERBB2, KIT, KRAS, MET, NRAS, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed; RNA analysis
81450	Targeted genomic sequence analysis panel, hematolymphoid neoplasm or disorder, 5-50 genes (eg, BRAF, CEBPA, DNMT3A, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MLL, NOTCH1, NPM1, NRAS), interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; DNA analysis or combined DNA and RNA analysis
81451	Targeted genomic sequence analysis panel, hematolymphoid neoplasm or disorder, 5-50 genes (eg, BRAF, CEBPA, DNMT3A, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MLL, NOTCH1, NPM1, NRAS), interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; RNA analysis
81455	Targeted genomic sequence analysis panel, solid organ or hematolymphoid neoplasm or disorder, 51 or greater genes (eg, ALK, BRAF, CDKN2A, CEBPA, DNMT3A, EGFR, ERBB2, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MET, MLL, NOTCH1, NPM1, NRAS, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; DNA analysis or combined DNA and RNA analysis

81456	Targeted genomic sequence analysis panel, solid organ or hematolymphoid neoplasm or disorder, 51 or greater genes (eg, ALK, BRAF, CDKN2A, CEBPA, DNMT3A, EGFR, ERBB2, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MET, MLL, NOTCH1, NPM1, NRAS, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; RNA analysis
81479	Unlisted molecular pathology procedure
81504	Oncology (tissue of origin), microarray gene expression profiling of > 2000 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as tissue similarity scores
81518	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 11 genes (7 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithms reported as percentage risk for metastatic recurrence and likelihood of benefit from extended endocrine therapy
81519	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 21 genes, utilizing formalin-fixed paraffin embedded tissue, algorithm reported as recurrence score
81520	Oncology (breast), mRNA gene expression profiling by hybrid capture of 58 genes (50 content and 8 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence risk score
81521	Oncology (breast), mRNA, microarray gene expression profiling of 70 content genes and 465 housekeeping genes, utilizing fresh frozen or formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk of distant metastasis
81522	Oncology (breast), mRNA, gene expression profiling by RT-PCR of 12 genes (8 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk score (Endopredict)
81523	Oncology (breast), mRNA, next-generation sequencing gene expression profiling of 70 content genes and 31 housekeeping genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk to distant metastasis - MAAA Breast Cancer Metastasis RNA Sequencing
81525	Oncology (colon), mRNA, gene expression profiling by real-time RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence score
81529	Oncology (cutaneous melanoma), mRNA, gene expression profiling by real-time RT-PCR of 31 genes (28 content and 3 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk, including likelihood of sentinel lymph node metastasis
81540	Oncology (tumor of unknown origin), mRNA, gene expression profiling by real-time RT-PCR of 92 genes (87 content and 5 housekeeping) to classify tumor into main cancer type and subtype, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported
81541	Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score
81542	Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score (Decipher)
81546	Oncology (thyroid), mRNA, gene expression analysis of 10,196 genes, utilizing fine needle aspirate, algorithm reported as a categorical result (eg, benign or suspicious)
81551	Oncology (prostate), promoter methylation profiling by real-time PCR of 3 genes (GSTP1, APC, RASSF1), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a likelihood of prostate cancer detection on repeat biopsy
81552	Oncology (uveal melanoma), mRNA, gene expression profiling by real-time RT-PCR of 15 genes (12 content and 3 housekeeping), utilizing fine needle aspirate or formalin-fixed paraffin-embedded tissue, algorithm reported as risk of metastasis
0005U	Oncology (prostate) gene expression profile by real-time RT-PCR of 3 genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score
0006M	Oncology (hepatic), mRNA expression levels of 161 genes, utilizing fresh hepatocellular carcinoma tumor tissue, with alpha-fetoprotein level, algorithm reported as a risk classifier
0007M	Oncology (gastrointestinal neuroendocrine tumors), real-time PCR expression analysis of 51 genes, utilizing whole peripheral blood, algorithm reported as a nomogram of tumor disease index
0011M	Oncology, prostate cancer, mRNA expression assay of 12 genes (10 content and 2 housekeeping), RT-PCR test utilizing blood plasma and/or urine, algorithms to predict high-grade prostate cancer risk
0012M	Oncology (urothelial), mRNA, gene expression profiling by real-time quantitative PCR of five genes (MDK, HOXA13, CDC2 [CDK1], IGFBP5, and XCR2), utilizing urine, algorithm reported as a risk score for having urothelial carcinoma
0013M	Oncology (urothelial), mRNA, gene expression profiling by real-time quantitative PCR of five genes (MDK, HOXA13, CDC2 [CDK1], IGFBP5, and CXCR2), utilizing urine, algorithm reported as a risk score for having recurrent urothelial carcinoma
0013U	Oncology (solid organ neoplasia), gene rearrangement detection by whole genome next-generation sequencing, DNA, fresh or frozen tissue or cells, report of specific gene rearrangement(s)
0014U	Hematology (hematolymphoid neoplasia), gene rearrangement detection by whole genome nextgeneration sequencing, DNA, whole blood or bone marrow, report of specific gene rearrangement(s)
0016M	Oncology (bladder), mRNA, microarray gene expression profiling of 219 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as molecular subtype (luminal, luminal infiltrated, basal, basal claudin-low, neuroendocrinelike)
0016U	Oncology (hematolymphoid neoplasia), RNA, BCR/ABL1 major and minor breakpoint fusion transcripts, quantitative PCR amplification, blood or bone marrow, report of fusion not detected or detected with quantitation

0017M	Oncology (diffuse large B-cell lymphoma [DLBCL]), mRNA, gene expression profiling by fluorescent probe hybridization of 20 genes, formalin-fixed paraffinembedded tissue, algorithm reported as cell of origin
0017U	Oncology (hematolymphoid neoplasia), JAK2 mutation, DNA, PCR amplification of exons 12-14 and sequence analysis, blood or bone marrow, report of JAK2 mutation not detected or detected
0018U	Oncology (thyroid), microRNA profiling by RT-PCR of 10 microRNA sequences, utilizing fine needle aspirate, algorithm reported as a positive or negative result for moderate to high risk of malignancy
0019U	Oncology, RNA, gene expression by whole transcriptome sequencing, formalin-fixed paraffin embedded tissue or fresh frozen tissue, predictive algorithm reported as potential targets for therapeutic agents
0022U	Targeted genomic sequence analysis panel, cholangiocarcinoma and non- small cell lung neoplasia, DNA and RNA analysis, 1 - 23 genes, interrogation for sequence variants and rearrangements, reported as presence/absence of variants and associated therapy(ies) to consider Targeted genomic sequence analysis panel, cholangiocarcinoma and non-small cell lung neoplasia, DNA and RNA analysis, 1-23 genes, interrogation for sequence variants and rearrangements, reported as presence/absence of variants and associated therapy(ies) to consider
0023U	Oncology (acute myelogenous leukemia), DNA, genotyping of internal tandem duplication, p.D835, p.I836, using mononuclear cells, reported as detection or nondetection of FLT3 mutation and indication for or against the use of midostaurin
0026U	Oncology (thyroid), DNA and mRNA of 112 genes, next-generation sequencing, fine needle aspirate of thyroid nodule, algorithmic analysis reported as a categorical result ("Positive, high probability of malignancy" or "Negative, low probability of malignancy")
0027U	JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) gene analysis, targeted sequence analysis exons 12-15
0036U	Exome (ie, somatic mutations), paired formalin-fixed paraffin-embedded tumor tissue and normal specimen, sequence analyses
0037U	Targeted genomic sequence analysis, solid organ neoplasm, DNA analysis of 324 genes, interrogation for sequence variants, gene copy number amplifications, gene rearrangements, microsatellite instability and tumor mutational burden
0040U	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis, major breakpoint, quantitative
0045U	Oncology (breast ductal carcinoma in situ), mRNA, gene expression profiling by realtime RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence score
0046U	FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia) internal tandem duplication (ITD) variants, quantitative
0047U	Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score
0048U	Oncology (solid organ neoplasia), DNA, targeted sequencing of protein-coding exons of 468 cancer-associated genes, including interrogation for somatic mutations and microsatellite instability, matched with normal specimens, utilizing formalin-fixed paraffin-embedded tumor tissue, report of clinically significant mutation(s)
0049U	NPM1 (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, quantitative
0050U	Targeted genomic sequence analysis panel, acute myelogenous leukemia, DNA analysis, 194 genes, interrogation for sequence variants, copy number variants or rearrangements
0056U	Hematology (acute myelogenous leukemia), DNA, whole genome nextgeneration sequencing to detect gene rearrangement(s), blood or bone marrow, report of specific gene rearrangement(s)
0069U	Oncology (colorectal), microRNA, RT-PCR expression profiling of miR-31-3p, formalin-fixed paraffin-embedded tissue, algorithm reported as an expression score
0089U	Oncology (melanoma), gene expression profiling by RTqPCR, PRAME and LINC00518, superficial collection using adhesive patch(es)
0090U	Oncology (cutaneous melanoma), mRNA gene expression profiling by RT-PCR of 23 genes (14 content and 9 housekeeping), utilizing formalin-fixed paraffin-embedded (FFPE) tissue, algorithm reported as a categorical result (ie, benign, intermediate, malignant) - myPath Melanoma
0111U	Oncology (colon cancer), targeted KRAS (codons 12, 13, and 61) and NRAS (codons 12, 13, and 61) gene analysis utilizing formalin-fixed paraffin-embedded tissue
0113U	Oncology (prostate), measurement of PCA3 and TMPRSS2-ERG in urine and PSA in serum following prostatic massage, by RNA amplification and fluorescencebased detection, algorithm reported as risk score
0114U	Gastroenterology (Barrett's esophagus), VIM and CCNA1 methylation analysis, esophageal cells, algorithm reported as likelihood for Barrett's esophagus
0120U	Oncology (B-cell lymphoma classification), mRNA, gene expression profiling by fluorescent probe hybridization of 58 genes (45 content and 13 housekeeping genes), formalin-fixed paraffin-embedded tissue, algorithm reported as likelihood for primary mediastinal B-cell lymphoma (PMBCL) and diffuse large B-cell lymphoma (DLBCL) with cell of origin subtyping in the latter
0153U	Oncology (breast), mRNA, gene expression profiling by next-generation sequencing of 101 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a triple negative breast cancer clinical subtype(s) with information on immune cell involvement
0154U	Oncology (urothelial cancer), RNA, analysis by real-time RT-PCR of the FGFR3 (fibroblast growth factor receptor 3) gene analysis (ie, p.R248C [c.742C>T], p.S249C [c.746C>G], p.G370C [c.1108G>T], p.Y373C [c.1118A>G], FGFR3-TACC3v1, and FGFR3-TACC3v3) utilizing formalin-fixed paraffin-embedded urothelial cancer tumor tissue, reported as FGFR gene alteration status

0155U	Oncology (breast cancer), DNA, PIK3CA (phosphatidylinositol-4,5-bisphosphate 3- kinase, catalytic subunit alpha) (eg, breast cancer) gene analysis (ie, p.C420R, p.E542K, p.E545A, p.E545D [g.1635G>T only], p.E545G, p.E545K, p.Q546E, p.Q546R, p.H1047L, p.H1047R, p.H1047Y), utilizing formalin-fixed paraffinembedded breast tumor tissue, reported as PIK3CA gene mutation status
0171U	Targeted genomic sequence analysis panel, acute myeloid leukemia, myelodysplastic syndrome, and myeloproliferative neoplasms, DNA analysis, 23 genes, interrogation for sequence variants, rearrangements and minimal residual disease, reported as presence/absence
0172U	Oncology (solid tumor as indicated by the label), somatic mutation analysis of BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) and analysis of homologous recombination deficiency pathways, DNA, formalin-fixed paraffin-embedded tissue, algorithm quantifying tumor genomic instability score
0177U	Oncology (breast cancer), DNA, PIK3CA (phosphatidylinositol-4,5-bisphosphate 3- kinase catalytic subunit alpha) gene analysis of 11 gene variants utilizing plasma, reported as PIK3CA gene mutation status
0179U	Oncology (non-small cell lung cancer), cell-free DNA, targeted sequence analysis of 23 genes (single nucleotide variations, insertions and deletions, fusions without prior knowledge of partner/breakpoint, copy number variations), with report of significant mutation(s)
0204U	Oncology (thyroid), mRNA, gene expression analysis of 593 genes (including BRAF, RAS, RET, PAX8, and NTRK) for sequence variants and rearrangements, utilizing fine needle aspirate, reported as detected or not detected
0208U	Oncology (medullary thyroid carcinoma), mRNA, gene expression analysis of 108 genes, utilizing fine needle aspirate, algorithm reported as positive or negative for medullary thyroid carcinoma
0211U	Oncology (pan-tumor), DNA and RNA by next-generation sequencing, utilizing formalin-fixed paraffin-embedded tissue, interpretative report for single nucleotide variants, copy number alterations, tumor mutational burden, and microsatellite instability, with therapy association
0228U	Oncology (prostate), multianalyte molecular profile by photometric detection of macromolecules adsorbed on nanosponge array slides with machine learning, utilizing first morning voided urine, algorithm reported as likelihood of prostate cancer
0229U	BCAT1 (Branched chain amino acid transaminase 1) and IKZF1 (IKAROS family zinc finger 1) (eg, colorectal cancer) promoter methylation analysis
0235U	PTEN (phosphatase and tensin homolog) (eg, Cowden syndrome, PTEN hamartoma tumor syndrome), full gene analysis, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions
0238U	Oncology (Lynch syndrome), genomic DNA sequence analysis of MLH1, MSH2, MSH6, PMS2, and EPCAM, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions
0239U	Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free DNA, analysis of 311 or more genes, interrogation for sequence variants, including substitutions, insertions, deletions, select rearrangements, and copy number variations
0242U	Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free circulating DNA analysis of 55-74 genes, interrogation for sequence variants, gene copy number amplifications, and gene rearrangements
0244U	Oncology (solid organ), DNA, comprehensive genomic profiling, 257 genes, interrogation for single-nucleotide variants, insertions/deletions, copy number alterations, gene rearrangements, tumor-mutational burden and microsatellite instability, utilizing formalin-fixed paraffinembedded tumor tissue
0245U	Oncology (thyroid), mutation analysis of 10 genes and 37 RNA fusions and expression of 4 mRNA markers using next-generation sequencing, fine needle aspirate, report includes associated risk of malignancy expressed as a percentage
0250U	Oncology (solid organ neoplasm), targeted genomic sequence DNA analysis of 505 genes, interrogation for somatic alterations (SNVs [single nucleotide variant], small insertions and deletions, one amplification, and four translocations), microsatellite instability and tumor-mutation burden
0262U	Oncology (solid tumor), gene expression profiling by real-time RT-PCR of 7 gene pathways (ER, AR, PI3K, MAPK, HH, TGFB, Notch), formalin-fixed paraffinembedded (FFPE), algorithm reported as gene pathway activity score
0285U	Oncology, response to radiation, cell-free DNA, quantitative branched chain DNA amplification, plasma, reported as a radiation toxicity score - RadTox™ cfDNA test
0287U	Oncology (thyroid), DNA and mRNA, nextgeneration sequencing analysis of 112 genes, fine needle aspirate or formalinixed paraffin-embedded (FFPE) tissue, algorithmic prediction of cancer recurrence, reported as a categorical risk result (low, intermediate, high) - ThyroSeq® CRC
0288U	Oncology (lung), mRNA, quantitative PCR analysis of 11 genes (BAG1, BRCA1, CDC6, CDK2AP1, ERBB3, FUT3, IL11, LCK, RND3, SH3BGR, WNT3A) and 3 reference genes (ESD, TBP, YAP1), formalin-fixed paraffin-embedded (FFPE) tumor tissue, algorithmic interpretation reported as a recurrence risk score - DetermaRx™
0296U	Oncology (oral and/or oropharyngeal cancer), gene expression profiling by RNA sequencing at least 20 molecular features (eg, human and/or microbial mRNA), saliva, algorithm reported as positive or negative for signature associated with malignancy - mRNA CancerDetect™
0297U	Oncology (pan tumor), whole genome sequencing of paired malignant and normal DNA specimens, fresh or formalinixed paraffin-embedded (FFPE) tissue, blood or bone marrow, comparative sequence analyses and variant identification - Praxis Somatic Whole Genome Sequencing

0298U	Oncology (pan tumor), whole transcriptome sequencing of paired malignant and normal RNA specimens, fresh or formalin-fixed paraffin-embedded (FFPE) tissue, blood or bone marrow, comparative sequence analyses and expression level and chimeric transcript identification - Praxis Somatic Transcriptome
0299U	Oncology (pan tumor), whole genome optical genome mapping of paired malignant and normal DNA specimens, fresh frozen tissue, blood, or bone marrow, comparative structural variant identification - Praxis Somatic Optical Genome Mapping
0300U	Oncology (pan tumor), whole genome sequencing and optical genome mapping of paired malignant and normal DNA specimens, fresh tissue, blood, or bone marrow, comparative sequence analyses and variant identification - Praxis Somatic Combined Whole Genome Sequencing and Optical Genome Mapping
0306U	Oncology (minimal residual disease [MRD]), next-generation targeted sequencing analysis, cell-free DNA, initial (baseline) assessment to determine a patient-specific panel for future comparisons to evaluate for MRD u(Do not report 0306U in conjunction with 0307U)t
0307U	Oncology (minimal residual disease [MRD]), next-generation targeted sequencing analysis of a patient-specific panel, cell-free DNA, subsequent assessment with comparison to previously analyzed patient specimens to evaluate for MRD(Do not report 0307U in conjunction with 0306U)
0313U	Oncology (pancreas), DNA and mRNA next-generation sequencing analysis of 74 genes and analysis of CEA (CEACAM5) gene expression, pancreatic cyst fluid, algorithm reported as a categorical result (ie, negative, low probability of neoplasia or positive, high probability of neoplasia)
0314U	Oncology (cutaneous melanoma), mRNA gene expression profiling by RT-PCR of 35 genes (32 content and 3 housekeeping), utilizing formalin-fixed paraffin-embedded (FFPE) tissue, algorithm reported as a categorical result (ie, benign, intermediate, malignant)
0315U	Oncology (cutaneous squamous cell carcinoma), mRNA gene expression profiling by RT-PCR of 40 genes (34 content and 6 housekeeping), utilizing formalin-fixed paraffinembedded tissue, algorithm reported as a categorical risk result (ie, Class 1, Class 2A, Class 2B)
0317U	Oncology (lung cancer), four-probe FISH (3q29, 3p22.1, 10q22.3, 10cen) assay, whole blood, predictive algorithmgenerated evaluation reported as decreased or increased risk for lung cancer
0326U	Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free circulating DNA analysis of 83 or more genes, interrogation for sequence variants, gene copy number amplifications, gene rearrangements, microsatellite instability and tumor mutational burden
0329U	Oncology (neoplasia), exome and transcriptome sequence analysis for sequence variants, gene copy number amplifications and deletions, gene rearrangements, microsatellite instability and tumor mutational burden utilizing DNA and RNA from tumor with DNA from normal blood or saliva for subtraction, report of clinically significant mutation(s) with therapy associations
0331U	Oncology (hematolymphoid neoplasia), optical genome mapping for copy number alterations and gene rearrangements utilizing DNA from blood or bone marrow, report of clinically significant alternations
0332U	Oncology (pan-tumor), genetic profiling of 8 DNA-regulatory (epigenetic) markers by quantitative polymerase chain reaction (qPCR), whole blood, reported as a high or low probability of responding to immune checkpoint–inhibitor therapy
0333U	Oncology (liver), surveillance for hepatocellular carcinoma (HCC) in highrisk patients, analysis of methylation patterns on circulating cell-free DNA (cfDNA) plus measurement of serum of AFP/AFP-L3 and oncoprotein des-gammacarboxy-prothrombin (DCP), algorithm reported as normal or abnormal result
0334U	Oncology (solid organ), targeted genomic sequence analysis, formalin-fixed paraffinembedded (FFPE) tumor tissue, DNA analysis, 84 or more genes, interrogation for sequence variants, gene copy number amplifications, gene rearrangements, microsatellite instability and tumor mutational burden
0339U	Oncology (prostate), mRNA expression profiling of HOXC6 and DLX1, reverse transcription polymerase chain reaction (RT-PCR), first-void urine following digital rectal examination, algorithm reported as probability of high-grade cancer
0340U	Oncology (pan-cancer), analysis of minimal residual disease (MRD) from plasma, with assays personalized to each patient based on prior next-generation sequencing of the patient’s tumor and germline DNA, reported as absence or presence of MRD, with disease-burden correlation, if appropriate
0343U	Oncology (prostate), exosome-based analysis of 442 small noncoding RNAs (sncRNAs) by quantitative reverse transcription polymerase chain reaction (RT-qPCR), urine, reported as molecular evidence of no-, low-, intermediate- or high-risk of prostate cancer
0356U	Oncology (oropharyngeal), evaluation of 17 DNA biomarkers using droplet digital PCR (ddPCR), cell-free DNA, algorithm reported as a prognostic risk score for cancer recurrence
0362U	Oncology (papillary thyroid cancer), gene-expression profiling via targeted hybrid capture–enrichment RNA sequencing of 82 content genes and 10 housekeeping genes, formalin-fixed paraffin embedded (FFPE) tissue, algorithm reported as one of three molecular subtypes
0363U	Oncology (urothelial), mRNA, geneexpression profiling by real-time quantitative PCR of 5 genes (MDK, HOXA13, CDC2 [CDK1], IGFBP5, and CXCR2), utilizing urine, algorithm incorporates age, sex, smoking history, and macrohematuria frequency, reported as a risk score for having urothelial carcinoma
G9840	KRAS gene mutation testing performed before initiation of anti-EGFR MoAb
G9841	KRAS gene mutation testing not performed before initiation of anti-EGFR MoAb
S3854	Gene expression profiling panel for use in the management of breast cancer treatment

History

Status	Review Date	Effective Date	Action
Created	09/21/2022	02/12/2023	Independent Multispecialty Physician Panel (IMPP) review. Original effective date.