



Status: Reaffirmed

Effective Date: 12/30/2023

Doc ID: REH02-1223.1-R1025

Last Review Date: 10/09/2025

Approval and implementation dates for specific health plans may vary. Please consult the applicable health plan for more details.

Clinical Appropriateness Guidelines

Outpatient Rehabilitative and Habilitative Services

Appropriate Use Criteria: Site of Care for Physical, Occupational, and Speech Therapies

Proprietary

© 2025 Carelon Medical Benefits Management, Inc. All rights reserved.

Table of Contents

Clinical Appropriateness Guidelines 1

Table of Contents 2

Description and Application of the Guidelines 3

General Clinical Guideline 4

 Clinical Appropriateness Framework 4

 Simultaneous Ordering of Multiple Diagnostic or Therapeutic Interventions 4

 Repeat Diagnostic Intervention 4

 Repeat Therapeutic Intervention 5

Site of Care for Physical, Occupational, and Speech Therapies 6

 General Information 6

 Clinical Indications 6

 Site of Care 6

 References 8

 Codes 8

History 9

Description and Application of the Guidelines

The Carelon Clinical Appropriateness Guidelines (hereinafter “the Carelon Clinical Appropriateness Guidelines” or the “Guidelines”) are designed to assist providers in making the most appropriate treatment decision for a specific clinical condition for an individual. The Guidelines establish objective and evidence-based criteria for medical necessity determinations, where possible, that can be used in support of the following:

- To establish criteria for when services are medically necessary
- To assist the practitioner as an educational tool
- To encourage standardization of medical practice patterns
- To curtail the performance of inappropriate and/or duplicate services
- To advocate for patient safety concerns
- To enhance the quality of health care
- To promote the most efficient and cost-effective use of services

The Carelon guideline development process complies with applicable accreditation and legal standards, including the requirement that the Guidelines be developed with involvement from appropriate providers with current clinical expertise relevant to the Guidelines under review and be based on the most up-to-date clinical principles and best practices. Resources reviewed include widely used treatment guidelines, randomized controlled trials or prospective cohort studies, and large systematic reviews or meta-analyses. Carelon reviews all of its Guidelines at least annually.

Carelon makes its Guidelines publicly available on its website. Copies of the Guidelines are also available upon oral or written request. Additional details, such as summaries of evidence, a list of the sources of evidence, and an explanation of the rationale that supports the adoption of the Guidelines, are included in each guideline document.

Although the Guidelines are publicly available, Carelon considers the Guidelines to be important, proprietary information of Carelon, which cannot be sold, assigned, leased, licensed, reproduced or distributed without the written consent of Carelon.

Carelon applies objective and evidence-based criteria, and takes individual circumstances and the local delivery system into account when determining the medical appropriateness of health care services. The Carelon Guidelines are just guidelines for the provision of specialty health services. These criteria are designed to guide both providers and reviewers to the most appropriate services based on a patient’s unique circumstances. In all cases, clinical judgment consistent with the standards of good medical practice should be used when applying the Guidelines. Guideline determinations are made based on the information provided at the time of the request. It is expected that medical necessity decisions may change as new information is provided or based on unique aspects of the patient’s condition. The treating clinician has final authority and responsibility for treatment decisions regarding the care of the patient and for justifying and demonstrating the existence of medical necessity for the requested service. The Guidelines are not a substitute for the experience and judgment of a physician or other health care professionals. Any clinician seeking to apply or consult the Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient’s care or treatment.

The Guidelines do not address coverage, benefit or other plan specific issues. Applicable federal and state coverage mandates take precedence over these clinical guidelines, and in the case of reviews for Medicare Advantage Plans, the Guidelines are only applied where there are not fully established CMS criteria. If requested by a health plan, Carelon will review requests based on health plan medical policy/guidelines in lieu of the Carelon Guidelines. Pharmaceuticals, radiotracers, or medical devices used in any of the diagnostic or therapeutic interventions listed in the Guidelines must be FDA approved or conditionally approved for the intended use. However, use of an FDA approved or conditionally approved product does not constitute medical necessity or guarantee reimbursement by the respective health plan.

The Guidelines may also be used by the health plan or by Carelon for purposes of provider education, or to review the medical necessity of services by any provider who has been notified of the need for medical necessity review, due to billing practices or claims that are not consistent with other providers in terms of frequency or some other manner.

General Clinical Guideline

Clinical Appropriateness Framework

Critical to any finding of clinical appropriateness under the guidelines for a specific diagnostic or therapeutic intervention are the following elements:

- Prior to any intervention, it is essential that the clinician confirm the diagnosis or establish its pretest likelihood based on a complete evaluation of the patient. This includes a history and physical examination and, where applicable, a review of relevant laboratory studies, diagnostic testing, and response to prior therapeutic intervention.
- The anticipated benefit of the recommended intervention should outweigh any potential harms including from delay or decreased access to services that may result (net benefit).
- Widely used treatment guidelines and/or current clinical literature and/or standards of medical practice should support that the recommended intervention offers the greatest net benefit among competing alternatives.
- There exists a reasonable likelihood that the intervention will change management and/or lead to an improved outcome for the patient.

Providers may be required to submit clinical documentation in support of a request for services. Such documentation must a) accurately reflect the clinical situation at the time of the requested service, and b) sufficiently document the ordering provider's clinical intent.

If these elements are not established with respect to a given request, the determination of appropriateness will most likely require a peer-to-peer conversation to understand the individual and unique facts that would justify a finding of clinical appropriateness. During the peer-to-peer conversation, factors such as patient acuity and setting of service may also be taken into account to the extent permitted by law.

Simultaneous Ordering of Multiple Diagnostic or Therapeutic Interventions

Requests for multiple diagnostic or therapeutic interventions at the same time will often require a peer-to-peer conversation to understand the individual circumstances that support the medical necessity of performing all interventions simultaneously. This is based on the fact that appropriateness of additional intervention is often dependent on the outcome of the initial intervention.

Additionally, either of the following may apply:

- Current literature and/or standards of medical practice support that one of the requested diagnostic or therapeutic interventions is more appropriate in the clinical situation presented; or
- One of the diagnostic or therapeutic interventions requested is more likely to improve patient outcomes based on current literature and/or standards of medical practice.

Repeat Diagnostic Intervention

In general, repeated testing of the same anatomic location for the same indication should be limited to evaluation following an intervention, or when there is a change in clinical status such that additional testing is required to determine next steps in management. At times, it may be necessary to repeat a test using different techniques or protocols to clarify a finding or result of the original study.

Repeated testing for the same indication using the same or similar technology may be subject to additional review or require peer-to-peer conversation in the following scenarios:

- Repeated diagnostic testing at the same facility due to technical issues
- Repeated diagnostic testing requested at a different facility due to provider preference or quality concerns

- Repeated diagnostic testing of the same anatomic area based on persistent symptoms with no clinical change, treatment, or intervention since the previous study
- Repeated diagnostic testing of the same anatomic area by different providers for the same member over a short period of time

Repeat Therapeutic Intervention

In general, repeated therapeutic intervention in the same anatomic area is considered appropriate when the prior intervention proved effective or beneficial and the expected duration of relief has lapsed. A repeat intervention requested prior to the expected duration of relief is not appropriate unless it can be confirmed that the prior intervention was never administered. Requests for ongoing services may depend on completion of previously authorized services in situations where a patient's response to authorized services is relevant to a determination of clinical appropriateness.

Site of Care for Physical, Occupational, and Speech Therapies

"Site of Care," "Site of Service" or another term such as "Setting" or "Place of Service" may be terms used in benefit plans, provider contracts, or other materials instead of or in addition to "Level of Care" and, in some plans, these terms may be used interchangeably.

General Information

Scope

These guidelines address site of care for outpatient physical (PT), occupational (OT), and speech therapies (ST) that are routinely performed outside of a hospital setting. Specifically, the guidelines outline the medical necessity criteria for providing such services at a hospital outpatient department (HOPD) where a higher level of care and support may be available relative to a freestanding or independent clinic. In short, the provision of therapies within HOPD facilities can be considered appropriate when they offer equipment and/or level of support which are unavailable at non-HOPD sites and necessary to render outpatient therapy services safely and effectively.

This guideline pertains to requests for outpatient rehabilitation and habilitation therapies for all disciplines (PT, OT, ST) for both pediatric and adult patients.

This guideline does not address the clinical appropriateness of individual procedures but rather restricts its focus to the selection of optimal site of care where the service is rendered. Some services may also be subject to clinical appropriateness review; in which case, a separate clinical guideline or policy may be used to adjudicate clinical appropriateness.*

This guideline does not address the availability of alternative non-hospital sites to perform an individual procedure. The purpose of this guideline is to define the clinical scenarios in which hospital-based care is medically necessary and, by exclusion, when it is clinically reasonable to provide services in a non-hospital setting.

See the [Codes](#) section for a list of services included in these guidelines.

**Note: Not all services that are subject to Site of Service review also require medical necessity review of the service itself. Some may only be subject to Site of Care review.*

Definitions

Hospital outpatient department (HOPD): Hospital-based department at which therapy services are rendered to outpatients

Non-hospital outpatient department (non-HOPD): Freestanding and/or off-campus therapy facility that is not operating as a HOPD

Clinical Indications

Site of Care

Providing PT, OT, or ST at a hospital outpatient site of care (HOPD) is considered medically necessary for requests that meet applicable medical necessity criteria for the service being performed in **ANY** of the following scenarios:

- Specialized equipment is necessary to achieve the therapeutic goals, which requires **BOTH** of the following:

- The specialized equipment is available only in the hospital-based facility (e.g., ceiling-mounted gait device that allows for variable weight-bearing, bariatric tables)
 - Therapeutic goals could not reasonably be achieved with use of alternative equipment that is more widely available
 - Specialized therapy and/or assessments (e.g., for certain equipment) are necessary to achieve therapeutic goals, which requires **ANY** of the following:
 - Subspecialized therapist (e.g., pediatric speech therapist)
 - Collaborative and coordinated care between more than one discipline (e.g., co-treatment with PT and OT, or therapist and wound team)
 - Clinical complexity requires access to higher level care, which requires **ANY** of the following:
 - Patient's underlying condition requires medical monitoring beyond the capabilities of the freestanding therapy facility
 - There is a potential need for immediate medical evaluation and care due to the patient's underlying condition and/or to the rendering of certain therapies (e.g., high risk of acute cardiac event, stroke, or seizures)
 - When performance of therapies outside the HOPD would reasonably be expected to create clinically significant delays in care
 - Geographic inaccessibility* of freestanding clinics presents a barrier to timely and appropriate therapy
- *Note: The definition of "inaccessibility" may vary per individual health plan policy.*

Rationale

With the widespread expansion of freestanding therapy clinics, access to routine therapy services in communities has greatly increased beyond the HOPD. As a result, overall reliance on HOPDs for therapies is less, which are more appropriately reserved for those situations in which the capabilities of HOPD therapy can uniquely improve care. While there are no studies formally comparing the quality of rehabilitative/habilitative therapies delivered in the HOPD and freestanding environments, there are some differentiating characteristics between the two settings which generally bear on determination of the most appropriate setting for a given clinical scenario. On a high level, this would include considerations related to equipment, medical needs, specialized care, and access. The need to perform HOPD-based therapies is therefore based on principle-based patient, personnel, or technical factors that are expected to offer net benefit to HOPD-based therapies by making them safer and/or more effective than therapies in freestanding facilities when available.

HOPD may offer specialized equipment and expertise that is not widely available in freestanding facilities. While the exact capabilities of freestanding facilities vary from market to market, the vast majority offer physical therapy and relatively few offer speech therapy, particularly pediatric speech therapy whose specialized practitioners typically practice only within a hospital (or school) setting. Similarly, HOPD therapy clinics are generally more likely to have advanced equipment (e.g., ceiling mounted gait-training device) and/or more extensive equipment to accommodate a broader array of patient populations (e.g., bariatric). However, wherein such equipment is available within a freestanding facility, or the use of such equipment is not essential to achieve the therapeutic goals, the medical necessity for a HOPD-based therapy is not established and delivering care in the freestanding facility would be appropriate.

Collaboration is another form of specialized care that is sometimes required to achieve the highest quality of care for a patient. This includes collaboration between therapy disciplines as well as collaboration between therapists and other providers. For instance, co-treatment of a traumatic brain injured patient by an occupational therapist and a speech therapist might be warranted to facilitate independence with self-care activities in the context of cognitive deficits. Additionally, that same patient's care might be more effective if his physical therapist is able to collaborate on-site with the physician who is medically managing his spasticity. Therefore, when such direct collaboration is needed and available only within the HOPD, the HOPD setting would be considered the most appropriate site.

Lastly, the primary principle guiding the site of care is ensuring the safety of the patient. Given the breadth of conditions and patients treated in outpatient therapy clinics, there is a broad range of medical needs, complexity, and risk. Risk for adverse events occurring during therapy can be due to the condition being treated (as in cardiac rehab), a related clinical feature (e.g., labile blood pressures associated with a spinal cord injury) or an unrelated comorbidity (e.g., angina in a knee replacement patient). In such situations, the potential need for proximal access to higher level care may warrant that therapy be rendered in a HOPD.

References

1. Bertelsen JB, Refsgaard J, Kanstrup H, et al. Hospital-based versus community-based shared care cardiac rehabilitation after acute coronary syndrome: protocol for a randomized clinical trial. *Danish Medical Journal*. 2013;60(9):A4699.
2. Fragala-Pinkham MA, Haley SM, Goodgold S. Evaluation of a community-based group fitness program for children with disabilities. *Pediatric Physical Therapy*. 2006;18(2):159-67.
3. American Physical Therapy Association. Access to, admission to, and patient/client rights within physical therapy services. Last updated August 30, 2018. Available at: <https://www.apta.org/apta-and-you/leadership-and-governance/policies/access-admission-patient-client-rights>. Accessed on November 14, 2022.
4. American Physical Therapy Association. Guide to physical therapist practice. 2nd Edition revised. American Physical Therapy Association. January 2003. Originally published as: Guide to Physical Therapist Practice. *Phys Ther*. 2001; 81:9-744.
5. American Physical Therapy Association. Physical therapists' role in prevention, wellness, fitness, health promotion, and management of disease and disability. September 20, 2019. Available at: <https://www.apta.org/apta-and-you/leadership-and-governance/policies/pt-role-advocacy>. Accessed on November 14, 2022.
6. American Physical Therapy Association. Physical therapy for older adults. Last updated September 20, 2019. Available at: <https://www.apta.org/apta-and-you/leadership-and-governance/policies/pt-for-older-adults>. Accessed on November 14, 2022.
7. American Physical Therapy Association. Standards of practice for physical therapy. Last updated August 12, 2020. Available at: <https://www.apta.org/apta-and-you/leadership-and-governance/policies/standards-of-practice-pt>. Accessed on November 14, 2022.
8. Centers for Medicare & Medicaid Services (CMS). Pub. 100-02, Chapter 15, Section 220. Coverage of Outpatient Rehabilitation Therapy Services (Physical Therapy, Occupational Therapy, and Speech-Language Pathology Services) Under Medical Insurance and Section 230 Practice of Physical Therapy, Occupational Therapy, and Speech-Language Pathology. July 12, 2019. Available at: <http://www.cms.hhs.gov/manuals/Downloads/bp102c15.pdf>. Accessed on November 14, 2022.
9. Centers for Medicare & Medicaid Services. National Coverage Determination for Institutional and Home Care Patient Education Programs. NCD#170.1. Effective date not posted. Available at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Accessed on November 14, 2022.
10. Hillegass E, Puthoff M, Frese EM, et al.; Guideline Development Group. Role of physical therapists in the management of individuals at risk for or diagnosed with venous thromboembolism: evidence-based clinical practice guideline. *Phys Ther*. 2016; 96(2):143-166.

Codes

The following code list is not meant to be all-inclusive. Authorization requirements will vary by health plan. Please consult the applicable health plan for guidance on specific procedure codes.

Specific CPT codes for services should be used when available. Nonspecific or not otherwise classified codes may be subject to additional documentation requirements and review.

CPT/HCPCS

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright by the American Medical Association. All Rights Reserved. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

See link below for CPT code list.

<https://providers.carelonmedicalbenefitsmanagement.com/rehabilitation/resources/>

ICD-10 Diagnosis

Refer to the ICD-10 CM manual

History

Status	Review Date	Effective Date	Action
Reaffirmed	10/09/2025	Unchanged	Independent Multispecialty Physician Panel (IMPP) review. Guideline reaffirmed.
Reaffirmed	04/15/2024	Unchanged	IMPP review. Scope and Rationale sections updated and aligned with all Carelon MBM Site of Care Guidelines. Expanded Site of Care guideline title for clarity. Added required language to General Clinical Guideline per new Medicare regulations.
Created	07/18/2023	12/30/2023	IMPP review. Original effective date.