Testicular (Germ Cell Tumors) Cancer Pathways

Patient Name:	Date of Birth:
Member Number:	Treatment Start Date:
Pathology:	Stage:
Line of Therapy:	ICD-10 Code:
Primary Therapy	
Good Risk	
Pure Seminoma (Stages IS-IIIB) and Non-Seminoma (Stages IS-IIIA)	
☐ BEP: bleomycin, etoposide, and cisplat	in*
□ EP: etoposide and cisplatin [†]	
Intermediate Risk	
Pure Seminoma (Stages IIC-IIIC) and Non-Seminoma (Stages IIIB-IIIC)	
□ BEP: bleomycin, etoposide, and cisplat	in*
VIP: etoposide, ifosfamide and cisplatin	
Poor Risk	
 Non-Seminoma (Stages IIIB-IIIC) 	
☐ BEP: bleomycin, etoposide, and cisplat	in
VIP: etoposide, ifosfamide and cisplatin	

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.



^{*} BEP is typically given for 3 cycles in good risk seminoma, and 4 cycles in intermediate risk

[†] EP given for 4 cycles